

# 1 Posets and their Hasse diagrams

**Definition I.** A *partially ordered set* (often abbreviated to *poset*)  $\mathcal{P} = (P, \leq)$  is a set  $P$  (called the *ground set*) with a *partial order*  $\leq$ . A partial order  $\leq$  is a *relation* on  $P$  (that is, a subset of  $P \times P$ ) that is

- *reflexive*:  $a \leq a$  for all  $a \in P$ .
- *antisymmetric*: if  $a \leq b$  and  $b \leq a$ , then  $a = b$ .
- *transitive*: if  $a \leq b$  and  $b \leq c$  then  $a \leq c$ .

If  $a \leq b$  then we say  $a$  is *less than*  $b$  or  $a$  *precedes*  $b$ . If  $a \leq b$  or  $b \leq a$ , we say  $a$  and  $b$  are *comparable*. Otherwise, they are *incomparable*. A poset is *totally ordered* if every pair of elements are comparable.

We write  $a < b$  if  $a \leq b$  and  $a \neq b$ .

We say that  $b$  *covers*  $a$  if  $a < b$  and there does not exist any third element  $x$  such that  $a < x < b$ .

A *subposet* of a poset  $(P, \leq)$  is a subset of  $P$  with the restriction of the partial order  $\leq$ .

A *chain* in a poset  $(P, \leq)$  is a totally ordered subposet  $a_0 < a_1 < \dots < a_p$ . The *height* of an element  $a$  in a poset is the maximum number  $p$  (possibly infinity) such that there exists a chain  $a_0 < a_1 < \dots < a_p = a$  of length  $(p+1)$ .

Given a poset  $(P, \leq)$  and  $a, b$  in  $P$ , the *interval*  $[a, b]$  (also denoted  $P_{[a,b]}$ ) is the subposet

$$P_{[a,b]} = \{x \in P \mid a \leq x \leq b\}.$$


We also define the following notation for the subposets

$$P_{<a} = \{x \in P \mid x < a\} \quad P_{\leq a} = \{x \in P \mid x \leq a\} \quad P_{\geq a} = \{x \in P \mid x \geq a\} \quad P_{>a} = \{x \in P \mid x > a\}.$$

**Definition II.** A *Hasse diagram* is a directed graph that encodes a finite poset  $(P, \leq)$ . It consists of a vertex for each element of  $P$ , with a directed edge (usually oriented upwards) from  $a \in P$  to  $b \in P$  whenever  $b$  covers  $a$ .

It is convenient to write elements of  $P$  of height  $h$  on row  $(h+1)$  from the bottom of the Hasse diagram.

**Example III.** The following figures show the Hasse diagram of the poset of subsets of a set  $\{a, b, c\}$ , ordered by inclusion, and the Hasse diagram of the poset of partitions of the set  $\{a, b, c, d\}$ , ordered by refinement, and the Hasse diagram of the poset of divisors of 60, ordered by divisibility.



**Exercise 1.** Consider the set  $[3] = \{1, 2, 3\}$  with the usual ordering  $1 < 2 < 3$ . Draw the Hasse diagrams for the following partial orders on the Cartesian product  $[3] \times [3]$ .

- (a) *Lexicographical order*:  $(a, b) \leq (c, d)$  if  $a \leq c$ , or if  $a = c$  and  $b \leq d$ .
- (b) *Product order*:  $(a, b) \leq (c, d)$  if  $a \leq c$  and  $b \leq d$ .
- (c) *Reflexive closure of strict direct product order*:  $(a \leq b) \leq (c, d)$  if  $(a, b) = (c, d)$ , or if  $a < c$  and  $b < d$ .

## 1.1 Least elements, minima, and lower bounds

**Definition IV.** An element  $a$  in a poset  $(P, \leq)$  is called *minimal* if there is no element  $x$  with  $x \leq a$ . It is called a *least element* if  $a \leq b$  for all  $b \in P$ . *Maximal* elements and *greatest elements* are defined analogously.

A least element, if it exists, is unique. However, posets with no least element may have multiple (incomparable) minimal elements.

**Definition V.** Let  $A$  be a subset of a poset  $(P, \leq)$ . A *lower bound* of  $A$  is an element  $\ell \in P$  such that  $\ell \leq a$  for all  $a \in A$ . The element  $\ell$  may or may not be contained in  $A$ . A greatest element of the subposet of lower bounds of  $A$  is called the *greatest lower bound* of  $A$ . *Upper bounds* and *least upper bounds* are defined similarly.

**Definition VI.** A poset  $(P, \leq)$  is called a *lattice* if every pair of elements  $\{a, b\} \subseteq P$  has a greatest lower bound (denoted  $a \wedge b$ ) and a least upper bound (denoted  $a \vee b$ ).


## 1.2 Maps of posets

**Definition VII.** Given posets  $(P, \leq)$  and  $(Q, \leq)$ , a function  $f : P \rightarrow Q$  is called *order-preserving* or *monotone* if  $f(a) \leq f(b)$  whenever  $a \leq b$ . It is called *strictly order-preserving* or *strictly monotone* if  $f(a) < f(b)$  whenever  $a < b$ .

## 2 The order complex of a poset

**Definition VIII.** Let  $\mathcal{P} = (P, \leq)$  be a poset. The *order complex*  $\Delta(\mathcal{P})$  of  $\mathcal{P}$  is the abstract simplicial complex whose vertex set is  $P$ , and whose simplices are precisely the nonempty finite chains in  $\mathcal{P}$ .

**Example IX.** Consider the poset of nonempty subsets of  $\{1, 2, 3\}$  ordered by inclusion. The Hasse diagram and its order complex are illustrated. The geometric realization of the order complex is the barycentric subdivision of a 2-simplex.



**Exercise 2.** Verify that a monotone map of posets induces a simplicial map on their order complexes.

**Definition X.** A *flag complex* is an (abstract) simplicial complex  $X$  with the following property. Let  $S$  be a nonempty subset of vertices of  $X$ . If every pair of vertices of  $S$  span an edge, then  $S$  is a simplex of  $X$ .

**Exercise 3.** Prove that the order complex of a poset is a flag complex.