References
for
Chebyshev and Fourier Spectral Methods

Second Edition

John P. Boyd

University of Michigan
Ann Arbor, Michigan 48109-2143
email: jpboyd@engin.umich.edu
http://www-personal.engin.umich.edu/~jpboyd/

2000

590

,

Bibliography

Abadi, M. H. A. and Ortiz, E. L.: 1991, A tau method based on non-uniform space-time elements for the numerical simulation of solitons, *Computers Math. Applic.* **22**(9), 7–19.

Abarbanel, S. and Gottlieb, D.: 1985, Information content in spectral calculations, *in* E. M. Murman and S. S. Abarbanel (eds), *Progress and Supercomputing in Computational Fluid Dynamics*, number 6 in *Proc. U. S.-Israel Workshop*, Birkhauser, Boston, pp. 345–356.

Abarbanel, S., Gottlieb, D. and Carpenter, M. H.: 1996, On the removal of boundary errors caused by Runge-Kutta integration of nonlinear partial differential equations, *SIAM Journal of Scientific Computing* **17**(3), 777–782. Not spectral, but essential in optimizing Runge-Kutta methods with Chebyshev or other high order spatial discretizations.

Abarbanel, S., Gottlieb, D. and Tadmor, E.: 1986, Spectral methods for discontinuous problems, *in* K. W. Morton and M. J. Baines (eds), *Numerical Methods for Fluid Dynamics, II*, Oxford University Press, Oxford, pp. 129–153.

Abe, K. and Inoue, O.: 1980, Fourier expansion solution of the Korteweg-deVries equation, *Journal of Computational Physics* **34**, 202–210.

Abramowitz, M. and Stegun, I. A.: 1965, *Handbook of Mathematical Functions*, Dover, New York.

Alpert, B. A. and Rokhlin, V.: 1991, A fast algorithm for the evaluation of Legendre expansions, *SIAM Journal of Scientific Computing* **12**, 158–179. Provides a fast Legendre-to-Chebyshev transform.

Amon, C. H. and Patera, A. T.: 1989, Numerical calculation of stable three-dimensional tertiary states in grooved-channel flow, *Physics of Fluids A* **1**, 2005–2009.

Anagnostou, G., Maday, Y. and Patera, A. T.: 1991, Numerical simulation of incompressible fluid flows, *Concurrency-Practice and Experience* **3**(6), 667–685. Review: high order operator-integration factors methods, sliding mesh spectral mortar-element, and parallel computing.

Anderson, C. and Dahleh, M. D.: 1996, Rapid computation of the discrete Fourier transform, *SIAM Journal of Scientific Computing* **17**, 913–919. Off-grid interpolation to irregularly spaced points by using local Taylor expansions at a cost of about 8 FFTs.

Anderson, D. L. T.: 1973, An ocean model using parabolic cylinder functions, World Meteorological Organization, Geneva. Model of equatorial ocean where Hermite functions are the linearized normal modes; discontinued.

Aoyagi, A.: 1995, Nonlinear leapfrog instability for Fornberg's pattern, *Journal of Computational Physics* **120**, 316–322. Aliasing instability.

Arakawa, A.: 1966, Computational design for long-term numerical integration of the equations of fluid motion: Two dimensional incompressible flow, *Journal of Computational Physics* **1**, 119–143. Energy-conserving finite differences schemes.

Armstrong, T. P.: 1967, Numerical studies of nonlinear Vlasov equation, *Physics of Fluids* **10**, 1269–1280. Fourier-Galerkin in the space dimension, Hermite-Galerkin in the velocity coordinate. Slow convergence of the Hermite series, a difficulty not fully overcome until variable scaling introduced in Holloway, 1996a,b.

Arnold, V. I.: 1983, *Geometrical Methods in the Theory of Ordinary Differential Equations*, Springer-Verlag, New York, chapter 17, pp. 110–111. The algorithm of "counting the noses" to approximate irrational numbers by rational fractions.

Augenbaum, J.: 1989, An adaptive pseudospectral method for discontinuous problems, *Applied Numerical Mathematics* **5**, 459–480.

Augenbaum, J. M.: 1990, Multidomain adaptive pseudospectral methods for acoustic wave propagation in discontinuous media, *in* D. Lee, A. Cakmak and R. Vichnevetsky (eds), *Computational Acoustics-Seismo-Ocean Acoustics and Modeling*, number 3 in *Computational Acoustics*, Elsevier/North-Holland, Amsterdam, pp. 19–40.

Augenbaum, J. M.: 1992, The pseudospectral method for limited-area elastic wave calculations, *in* W. E. Fitzgibbon and M. F. Wheeler (eds), *Computational Methods in Geosciences*, Frontiers in Applied Mathematics, SIAM, Philadelphia. Chebyshev method needs only about half the resolution in each coordinate compared to fourth order differences.

Averbuch, A., Ioffe, L., Israeli, M. and Vozovoi, L.: 1998a, Two-dimensional parallel solver for the solution of navier-stokes equations with constant and variable coefficients using ADI on cells, *Parallel Comput.* **24**(5–6), 673–699. Local Fourier basis for nonperiodic problems, here with domain decomposition.

Averbuch, A., Israeli, M. and Vozovoi, L.: 1995, Parallel implementation of nonlinear evolution problems using parabolic domain decomposition, *Parallel Comput.* **21**(7), 1151–1183. Fourier method for nonperiodic boundary conditions; multidomain.

Averbuch, A., Israeli, M. and Vozovoi, L.: 1998b, A fast poisson solver of arbitrary order accuracy in rectangular regions, *SIAM J. Sci. Comput.* **19**(3), 933–952. Fourier method for nonperiodic boundary conditions with subtractions to handle Gibbs' Phenomenon and corner singularities.

Averbuch, A., Vozovoi, L. and Israeli, M.: 1997, On a fast direct elliptic solver by a modified fourier method, *Numer. Algorithms* **15**(3–4), 287–313. Fourier method for nonperiodic boundary conditions with subtractions to handle Gibbs' Phenomenon and corner singularities.

Babolian, E. and Delves, L. M.: 1979, An augmented Galerkin method for first kind Fredholm equations, *Journal of the Institute for Mathematics and its Applications* **24**, 151–174. Galerkin basis is Chebyshev polynomials.

Babolian, E. and Delves, L. M.: 1981, A fast Galerkin method for linear integro-differential equations, *IMA Journal of Numerical Analysis* **1**, 193–213. Chebyshev series.

Baer, F.: 1977, Adjustment of initial conditions required to suppress gravity oscillations in non-linear flows, *Contributions to Atmospheric Physics* **50**, 350–366.

Baer, F. and Tribbia, J. J.: 1977, On complete filtering of gravity modes through nonlinear initialization, *Monthly Weather Review* **105**, 1536–1539.

Bain, M.: 1978, Convergence theorems for Hermite functions and Jacobi polynomials, *Journal of the Institute for Mathematics and its Applications* **21**, 379–386.

Baker, Jr., G. A.: 1975, *Essentials of Padé Approximants*, Academic Press, New York. 220 pp.

Baker, M. D., Süli, E. and Ware, A. F.: 1992, Stability and convergence of the spectral Lagrange-Galerkin method for periodic and non-periodic convection-dominated diffusion problems, *Technical Report 92/19*, Oxford University Computing Laboratory, Numerical Analysis Group,Wolfson Bldg., Parks Rd., Oxford, England OX1 3QD.

Banerjee, K.: 1978, Hermite function solution of quantum anharmonic oscillator, *Proceedings Of The Royal Society of London, Series A* **364**, 264.

Banerjee, K., Bhatnagar, S. P., Choudhury, V. and Kanwal, S. S.: 1978, *Proceedings of the Royal Society of London A*. Hermite functions $\psi_n(\alpha y)$ with empirically-optimized scale factor α.

Bank, R. E. and Chan, T. F.: 1986, PLTMGC: A multi-grid continuation program for parameterized nonlinear elliptic systems, *SIAM Journal of Scientific and Statistical Computing* **7**, 540–559.

Barrett, E. W.: 1958, Eccentric circumpolar vortices in a barotropic atmosphere, *Tellus* **10**, 395–400. Spherical harmonics.

Basdevant, C., Deville, M., Haldenvvang, P., Lacroix, J., Orlandi, D., Patera, A., Peyret, R. and Quazzani, J.: 1986, Spectral and finite difference solutions of Burgers' equation, *Comput. Fluids* **14**, 23–41.

Bates, J. R.: 1984, An efficient semi-Lagrangian and alternating-direction implicit method for integrating the shallow-water equations, *Monthly Weather Review* **112**, 2033–2047.

Bates, J. R. and McDonald, A.: 1982, Multiply-upstream, semi-Lagrangian advective schemes: Analysis and application to a multilevel primitive equation model, *Monthly Weather Review* **110**, 1831–1842.

Bates, J. R., Semazzi, F. H. M., Higgins, R. W. and Barros, S. R. M.: 1990, Integration of the shallow-water equations on the sphere using a vector semi-Lagrangian scheme with a multigrid solver, *Monthly Weather Review* **118**, 1615–1627.

Baumgardner, J. R. and Frederickson, P. O.: 1985, Icosahedral discretization of the two-sphere, *SIAM Journal of Numerical Analysis* **22**, 1107–1115.

Bayliss, A. and Matkowsky, B.: 1987, Fronts, relaxation oscillations, and period-doubling in solid fuel combustion, *Journal of Computational Physics* **71**, 147–168. Adaptive pseudospectral grid.

Bayliss, A. and Matkowsky, B. J.: 1992, Nonlinear dynamics of cellular flames, *SIAM Journal of Applied Mathematics* **52**(2), 396–415. Chebyshev and Fourier pseudospectral combined with adaptive shifted arctan map and first order splitting time integration.

Bayliss, A. and Turkel, E.: 1992, Mappings and accuracy for Chebyshev pseudospectral approximations, *Journal of Computational Physics* **101**, 349–359.

Bayliss, A., Belytschko, T., Kulkarni, M. and Lott-Crumpler, D. A.: 1994, On the dynamics and role of imperfections for localization in thermo-viscoplastic materials, *Modelling and Simulation of Materials in Science and Engineering* **2**, 941–964. Adaptive Chebyshev pseudospectral method to compute adiabatic shear bands via time-marching with shifted arctan mapping.

Bayliss, A., Garbey, M. and Matkowsky, B. J.: 1995, Adaptive pseudo-spectral domain decomposition and the approximation of multiple layers, *Journal of Computational Physics* **119**, 132–141.

Bayliss, A., Gottlieb, D., matkowsky, B. J. and Minkoff, M.: 1989, An adaptive pseudo-spectral method for reaction-diffusion problems, *Journal of Computational Physics* **81**, 421–443.

Bayliss, A., Kuske, R. and Matkowsky, B. J.: 1990, A two-dimensional adaptive pseudo-spectral method, *Journal of Computational Physics* **91**, 174–196. Chebyshev polynomials and one-dimensional arctan/tan mapping with map parameters that are allowed to vary with the other spatial coordinate.

Belousov, S. L.: 1962, *Tables of Normalized Associate Legendre Polynomials*, Macmillan, New York.

Ben Belgacem, F. and Maday, Y.: 1997, The mortar element method for three dimensional finite elements, *Rairo-Math. Model Numer.* **31**(2), 289–302.

Bender, C. M. and Orszag, S. A.: 1978, *Advanced Mathematical Methods for Scientists and Engineers*, McGraw-Hill, New York. 594 pp.

Bermejo, R.: 1990, On the equivalence of semi-Lagrangian and particle-in-cell finite-element methods, *Monthly Weather Review* **118**, 979–987. Not spectral.

Bermejo, R. and Staniforth, A.: 1992, The conversion of semi-Lagrangian advection schemes to quasi-monotone schemes, *Monthly Weather Review* **120**, 2622–2632.

Bernardi, C. and Maday, Y.: 1992, *Approximations spectrales de problémes aux limites elliptiques*, Springer-Verlag France, Paris, France. 250 pp. In French; highly mathematical.

Bert, C. W. and Malik, M.: 1996, Differential quadrature method in computational mechanics: A review, *Applied Mechanics Reviews* **49**, 1–28. Pseudospectral method implemented as N-th order finite difference scheme.

Bialecki, B.: 1989, Sinc-type approximations in h^1 norm with application to boundary value problems, *Journal of Computational and Applied Mathematics* **25**, 289–303.

Bialecki, B.: 1991, Sinc-collocation methods for two-point boundary value problems, *IMA Journal of Numerical Analysis* **11**, 357–375.

Birkhoff, G. and Fix, G.: 1970, Variational methods for eigenvalue problems, *Numerical Solution of Field Problems in Continuum Mechanics*, Vol. 2, SIAM-AMS Proceedings of the American Mathematical Society, Providence, pp. 111–151. Hermite function and Fourier solution of eigenvalue problems.

Birkhoff, G. and Lynch, R. E.: 1984, *Numerical Solution of Elliptic Problems*, Society for Industrial and Applied Mathematics, Philadelphia. 300 pp.

Black, K.: 1998, Spectral elements on infinite domains, *SIAM J. Sci. Comput.* **19**(5), 1667–1681. Employs the change-of-coordinate of Boyd(1987b).

Blackburn, H. and Henderson, R. D.: 1996, Lock-in behavior in simulation vortex-induced simulation, *Experimental Thermal and Fluid Science* **12**, 184–189. Spectral element model of vibrations of a circular cylinder, induced by vibrations due to vortices in the surrounding fluid, using stiffly stable (Backward Differentiation) time integration. The grid is fixed to the (moving) cylinder and equations solved in this accelerating reference frame. ODEs for the motion of the cylinder are solved simultaneously with the hydrodynamic PDEs.

Blaisdell, G. A., T.Spyropoulos, E. and Qin, J. H.: 1996, The effect of the formulation of nonlinear terms on aliasing errors in spectral methods, *Applied Numerical Mathematics* **21**, 207–219. Show through both Burgers' equation and turbulence models that the skew-symmetric form of advection is more robust than the alternatives.

Blue, J. L. and Gummel, H. K.: 1970, Rational approximations to matrix exponential for systems of stiff differential equations, *Journal of Computational Physics* **5**, 70–83. Not spectral.

Boer, G. J., McFarlane, N. A., Laprise, R., Henderson, J. D. and Blanchet, J.-P.: 1984, The Canadian Climate Centre spectral atmospheric general circulation model, *Atmosphere-Ocean* **22**, 397–429.

Boffi, D. and Funaro, D.: 1994, An alternative approach to the analysis and the approximation of the Navier-Stokes equations, *Journal of Scientific Computing* **9**(1), 16.

Boomkamp, P. A. M., Boersma, B. J., Miesen, R. H. M. and Beijnon, G. V.: 1997, A Chebyshev collocation method for solving two-phase flow stability problems, *J. Comput. Phys.* **132**, 191–200. Collocation/QZ algorithm for eigenvalue problems, using a three-subdomain strategy.

Böttcher, M.: 1996, A semi-Lagrangian advection scheme with modified exponential splines, *Monthly Weather Review* **124**, 716. Not spectral.

Bouaoudi and Marcus, P. S.: 1991, Fast and accurate spectral treatment of coordinate singularities, *Journal of Computational Physics* **96**, 217–223. Polar and spherical coordinates using Robert functions. In later work, the authors abandoned this approach because this basis is highly ill-conditioned.

Boukir, K., Maday, Y. and Métivet, B.: 1997, A high order characteristics method for the incompressible Navier-Stokes equations, *Internat. J. Numer. Meth. Fluids* **25**(12), 1421–1454.

Bourke, W.: 1972, An efficient, one-level primitive-equation spectral model, *Monthly Weather Review* **100**, 683–689. Spherical harmonics.

Bourke, W.: 1974, A multi-level spectral model. I. Formulation and hemispheric integrations, *Monthly Weather Review* **102**, 687–701.

Bourke, W., McAvaney, K., Puri, K. and Thurling, R.: 1977, *Global modelling of atmospheric flow by spectral methods*, Vol. 17 of *Methods in Computational Physics*, Academic Press, New York, pp. 267–334.

Bouteloup, Y.: 1995, Improvement of the spectral representation of the earth topography, *Mon. Wea. Rev.* **123**, 1560–1573.

Bowers, K. L. and Lund, J.: 1987, Numerical solution of singular Poisson equations by the sinc-Galerkin method, *SIAM Journal of Numerical Analysis* **24**, 36–51.

Boyd, J. P.: 1978a, A Chebyshev polynomial method for computing analytic solutions to eigenvalue problems with application to the anharmonic oscillator, *Journal of Mathematical Physics* **19**, 1445–1456.

Boyd, J. P.: 1978b, The choice of spectral functions on a sphere for boundary and eigenvalue problems: A comparison of Chebyshev, Fourier and associated Legendre expansions, *Monthly Weather Review* **106**, 1184–1191.

Boyd, J. P.: 1978c, Spectral and pseudospectral methods for eigenvalue and nonseparable boundary value problems, *Monthly Weather Review* **106**, 1192–1203.

Boyd, J. P.: 1980a, The nonlinear equatorial Kelvin wave, *Journal of Physical Oceanography* **10**, 1–11.

Boyd, J. P.: 1980b, The rate of convergence of Hermite function series, *Mathematics of Computation* **35**, 1309–1316.

Boyd, J. P.: 1980c, Equatorial solitary waves, Part I: Rossby solitons, *Journal of Physical Oceanography* **10**, 1699–1718.

Boyd, J. P.: 1981a, A Sturm-Liouville eigenproblem with an interior pole, *Journal of Physical Oceanography* **22**, 1575–1590. Background on waves with critical points; nothing on spectral methods.

Boyd, J. P.: 1981b, The rate of convergence of Chebyshev polynomials for functions which have asymptotic power series about one endpoint, *Journal of Physical Oceanography* **37**, 189–196.

Boyd, J. P.: 1981c, The Moonbow, *Isaac Asimov's SF Mag.* **5**, 18–37. Properties of a toroidal world.

Boyd, J. P.: 1982a, The optimization of convergence for Chebyshev polynomial methods in an unbounded domain, *Journal of Computational Physics* **45**, 43–79. Infinite and semi-infinite intervals; guidelines for choosing the map parameter or domain size L.

Boyd, J. P.: 1982b, The effects of meridional shear on planetary waves, Part I: Nonsingular profiles, *Journal of the Atmospheric Sciences* **39**, 756–769.

Boyd, J. P.: 1982c, The effects of meridional shear on planetary waves, Part II: Critical latitudes, *Journal of the Atmospheric Sciences* **39**, 770–790. First application of cubic-plus-linear mapping with spectral methods. The detour procedure of Boyd (1985a) is better in this context.

Boyd, J. P.: 1982d, A Chebyshev polynomial rate-of-convergence theorem for Stieltjes functions, *Mathematics of Computation* **39**, 201–206.

Boyd, J. P.: 1982e, Theta functions, Gaussian series, and spatially periodic solutions of the Korteweg-de Vries equation, *Journal of Mathematical Physics* **23**, 375–387.

Boyd, J. P.: 1983a, Equatorial solitary waves, Part II: Envelope solitons, *Journal of Physical Oceanography* **13**, 428–449. This and the next two papers use Hermite series to solve linear, separable equations in perturbation theory for nonlinear waves.

Boyd, J. P.: 1983b, Long wave/short wave resonance in equatorial waves, *Journal of Physical Oceanography* **13**, 450–458.

Boyd, J. P.: 1983c, Second harmonic resonance for equatorial waves, *Journal of Physical Oceanography* **13**, 459–466.

Boyd, J. P.: 1983d, The continuous spectrum of linear Couette flow with the beta effect, *Journal of the Atmospheric Sciences* **40**, 2304–2308.

Boyd, J. P.: 1984a, The asymptotic coefficients of Hermite series, *Journal of Computational Physics* **54**, 382–410.

Boyd, J. P.: 1984b, Equatorial solitary waves, Part IV: Kelvin solitons in a shear flow, *Dynamics of Atmospheres and Oceans* **8**, 173–184.

Boyd, J. P.: 1984c, Cnoidal waves as exact sums of repeated solitary waves: New series for elliptic functions, *SIAM Journal of Applied Mathematics* **44**, 952–955. Imbricate series for nonlinear waves.

Boyd, J. P.: 1984d, The double cnoidal wave of the Korteweg-de Vries equation: An overview, *Journal of the Mathematical Physics* **25**, 3390–3401.

Boyd, J. P.: 1984e, Perturbation theory for the double cnoidal wave of the Korteweg-de Vries equation, *Journal of the Mathematical Physics* **25**, 3402–3414.

Boyd, J. P.: 1984f, The special modular transformation for the polycnoidal waves of the Korteweg-de Vries equation, *Journal of the Mathematical Physics* **25**, 3390–3401.

Boyd, J. P.: 1984g, Earthflight, *in* S. Shwarz (ed.), *Habitats*, DAW, New York, pp. 201–218. Toroidal planet.

Boyd, J. P.: 1985a, Complex coordinate methods for hydrodynamic instabilities and Sturm-Liouville problems with an interior singularity, *Journal of Computational Physics* **57**, 454–471.

Boyd, J. P.: 1985b, Equatorial solitary waves, Part 3: Modons, *Journal of Physical Oceanography* **15**, 46–54.

Boyd, J. P.: 1985c, An analytical and numerical study of the two-dimensional Bratu equation, *Journal of Scientific Computing* **1**, 183–206. Nonlinear eigenvalue problem with 8-fold symmetry.

Boyd, J. P.: 1985d, Barotropic equatorial waves: The non-uniformity of the equatorial beta-plane, *Journal of the Atmospheric Sciences* **42**, 1965–1967.

Boyd, J. P.: 1986a, Solitons from sine waves: analytical and numerical methods for non-integrable solitary and cnoidal waves, *Physica D* **21**, 227–246. Fourier pseudospectral with continuation and the Newton-Kantorovich iteration.

Boyd, J. P.: 1986b, Polynomial series versus sinc expansions for functions with corner or endpoint singularities, *Journal of Computational Physics* **64**, 266–269.

Boyd, J. P.: 1987a, Exponentially convergent Fourier/Chebyshev quadrature schemes on bounded and infinite intervals, *Journal of Scientific Computing* **2**, 99–109.

Boyd, J. P.: 1987b, Spectral methods using rational basis functions on an infinite interval, *Journal of Computational Physics* **69**, 112–142.

Boyd, J. P.: 1987c, Orthogonal rational functions on a semi-infinite interval, *Journal of Computational Physics* **70**, 63–88.

Boyd, J. P.: 1987d, Generalized solitary and cnoidal waves, *in* G. Brantstator, J. J. Tribbia and R. Madden (eds), *NCAR Colloquium on Low Frequency Variability in the Atmosphere*, National Center for Atmospheric Research, Boulder, Colorado, pp. 717–722. Numerical calculation of the exponentially small wings of the ϕ^4 breather.

Boyd, J. P.: 1988a, Chebyshev domain truncation is inferior to Fourier domain truncation for solving problems on an infinite interval, *Journal of Scientific Computing* **3**, 109–120.

Boyd, J. P.: 1988b, An analytical solution for a nonlinear differential equation with logarithmic decay, *Advances in Applied Mathematics* **9**, 358–363.

Boyd, J. P.: 1988c, The superiority of Fourier domain truncation to Chebyshev domain truncation for solving problems on an infinite interval, *Journal of Scientific Computing* **3**, 109–120.

Boyd, J. P.: 1989a, *Chebyshev and Fourier Spectral Methods*, Springer-Verlag, New York. 792 pp.

Boyd, J. P.: 1989b, New directions in solitons and nonlinear periodic waves: Polycnoidal waves, imbricated solitons, weakly non-local solitary waves and numerical boundary value algorithms, *in* T.-Y. Wu and J. W. Hutchinson (eds), *Advances in Applied Mechanics*, number 27 in *Advances in Applied Mechanics*, Academic Press, New York, pp. 1–82.

Boyd, J. P.: 1989c, Periodic solutions generated by superposition of solitary waves for the quarticly nonlinear Korteweg-de Vries equation, *ZAMP* **40**, 940–944. Imbrication of solitary wave generates good *approximate* periodic solutions.

Boyd, J. P.: 1989d, The asymptotic Chebyshev coefficients for functions with logarithmic endpoint singularities, *Applied Mathematics and Computation* **29**, 49–67.

Boyd, J. P.: 1989e, Non-local equatorial solitary waves, *in* J. C. J. Nihoul and B. M. Jamart (eds), *Mesoscale/Synoptic Coherent Structures in Geophysical Turbulence: Proc. 20th Liege Coll. on Hydrodynamics*, Elsevier, Amsterdam, pp. 103–112.

Boyd, J. P.: 1990a, The orthogonal rational functions of Higgins and Christov and Chebyshev polynomials, *Journal of Approximation Theory* **61**, 98–103.

Boyd, J. P.: 1990b, A numerical calculation of a weakly non-local solitary wave: the ϕ^4 breather, *Nonlinearity* **3**, 177–195.

Boyd, J. P.: 1990c, The envelope of the error for Chebyshev and Fourier interpolation, *Journal of Scientific Computing* **5**, 311–363.

Boyd, J. P.: 1990d, A Chebyshev/radiation function pseudospectral method for wave scattering, *Computers in Physics* **4**, 83–85.

Boyd, J. P.: 1991a, A comparison of numerical and analytical methods for the reduced wave equation with multiple spatial scales, *Applied Numerical Mathematics* **7**, 453–479.

Boyd, J. P.: 1991b, Monopolar and dipolar vortex solitons in two space dimensions, *Wave Motion* **57**, 223–243.

Boyd, J. P.: 1991c, Nonlinear equatorial waves, *in* A. R. Osborne (ed.), *Nonlinear Topics of Ocean Physics: Fermi Summer School, Course LIX*, North-Holland, Amsterdam, pp. 51–97.

Boyd, J. P.: 1991d, Weakly nonlocal solitary waves, *in* A. R. Osborne (ed.), *Nonlinear Topics of Ocean Physics: Fermi Summer School, Course LIX*, North-Holland, Amsterdam, pp. 527–556.

Boyd, J. P.: 1991e, Weakly nonlocal solitons for capillary-gravity waves: Fifth-degree Korteweg-de Vries equation, *Physica D* **48**, 129–146.

Boyd, J. P.: 1991f, Sum-accelerated pseudospectral methods: The Euler-accelerated sinc algorithm, *Applied Numerical Mathematics* **7**, 287–296.

Boyd, J. P.: 1992a, The arctan/tan and Kepler-Burger mappings for periodic solutions with a shock, front, or internal boundary layer, *Journal of Computational Physics* **98**, 181–193. Numerical trick which is useful for solitary waves and cnoidal waves.

Boyd, J. P.: 1992b, The energy spectrum of fronts: The time evolution of shocks in Burgers' equation, *Journal of the Atmospheric Sciences* **49**, 128–139.

Boyd, J. P.: 1992c, Multipole expansions and pseudospectral cardinal functions: A new generalization of the Fast Fourier Transform, *Journal of Computational Physics* **102**, 184–186.

Boyd, J. P.: 1992d, A fast algorithm for Chebyshev and Fourier interpolation onto an irregular grid, *Journal of Computational Physics* **103**, 243–257.

Boyd, J. P.: 1992e, Defeating the Runge phenomenon for equispaced polynomial interpolation via Tikhonov regularization, *Applied Mathematics Letters* **5**, 57–59.

Boyd, J. P.: 1993, Chebyshev and Legendre spectral methods in algebraic manipulation languages, *Journal of Symbolic Computing* **16**, 377–399.

Boyd, J. P.: 1994a, Hyperviscous shock layers and diffusion zones: Monotonicity, spectral viscosity, and pseudospectral methods for high order differential equations, *Journal of Scientific Computing* **9**, 81–106.

Boyd, J. P.: 1994b, The rate of convergence of Fourier coefficients for entire functions of infinite order with application to the Weideman-Cloot sinh-mapping for pseudospectral computations on an infinite interval, *Journal of Computational Physics* **110**, 360–372.

Boyd, J. P.: 1994c, The slow manifold of a five mode model, *Journal of the Atmospheric Sciences* **51**, 1057–1064.

Boyd, J. P.: 1994d, Nonlocal modons on the beta-plane, *Geophysical and Astrophysical Fluid Dynamics* **75**, 163–182.

Boyd, J. P.: 1994e, Time-marching on the slow manifold: The relationship between the nonlinear Galerkin method and implicit timestepping algorithms, *Applied Mathematics Letters* **7**, 95–99.

Boyd, J. P.: 1994f, Sum-accelerated pseudospectral methods: Finite differences and sech-weighted differences, *Computer Methods in Applied Mechanics and Engineering* **116**, 1–11.

Boyd, J. P.: 1995a, Weakly nonlocal envelope solitary waves: Numerical calculations for the Klein-Gordon (ϕ^4) equation, *Wave Motion* **21**, 311–330.

Boyd, J. P.: 1995b, A hyperasymptotic perturbative method for computing the radiation coefficient for weakly nonlocal solitary waves, *Journal of Computational Physics* **120**, 15–32.

Boyd, J. P.: 1995c, Eight definitions of the slow manifold: Seiches, pseudoseiches and exponential smallness, *Dynamics of Atmospheres and Oceans* **22**, 49–75.

Boyd, J. P.: 1995d, A lag-averaged generalization of Euler's method for accelerating series, *Applied Mathematics and Computation* **72**, 146–166.

Boyd, J. P.: 1995e, A Chebyshev polynomial interval-searching method ("Lanczos economization") for solving a nonlinear equation with application to the nonlinear eigenvalue problem, *Journal of Computational Physics* **118**, 1–8.

Boyd, J. P.: 1995f, Multiple precision pseudospectral computations of the radiation coefficient for weakly nonlocal solitary waves: Fifth-Order Korteweg-deVries equation, *Computers in Physics* **9**, 324–334.

Boyd, J. P.: 1996a, Asymptotic Chebyshev coefficients for two functions with very rapidly or very slowly divergent power series about one endpoint, *Applied Mathematics Letters* **9**(2), 11–15.

Boyd, J. P.: 1996b, Traps and snares in eigenvalue calculations with application to pseudospectral computations of ocean tides in a basin bounded by meridians, *Journal of Computational Physics* **126**, 11–20. Corrigendum, **136**, no. 1, 227-228 (1997).

Boyd, J. P.: 1996c, Numerical computations of a nearly singular nonlinear equation: Weakly nonlocal bound states of solitons for the Fifth-Order Korteweg-deVries equation, *Journal of Computational Physics* **124**, 55–70.

Boyd, J. P.: 1996d, The Erfc-Log filter and the asymptotics of the Vandeven and Euler sequence accelerations, *in* A. V. Ilin and L. R. Scott (eds), *Proceedings of the Third International Conference on Spectral and High Order Methods*, Houston Journal of Mathematics, Houston, Texas, pp. 267–276.

Boyd, J. P.: 1997a, Padé approximant algorithm for solving nonlinear ODE boundary value broblems on an unbounded domain, *Computers and Physics* **11**(3), 299–303.

Boyd, J. P.: 1997b, Pseudospectral/Delves-Freeman computations of the radiation coefficient for weakly nonlocal solitary waves of the Third Order Nonlinear Schroedinger Equation and their relation to hyperasymptotic perturbation theory, *Journal of Computational Physics* **138**, 665–694.

Boyd, J. P.: 1997c, The periodic generalization of Camassa-Holm "peakons": An exact superposition of solitary waves, *Applied Mathematics and Computation* **81**(2), 173–187. Classical solitons.

Boyd, J. P.: 1997d, Construction of Lighthill's unitary functions: The imbricate series of unity, *Applied Mathematics and Computation* **86**, 1–10.

Boyd, J. P.: 1998a, Radiative decay of weakly nonlocal solitary waves, *Wave Motion* **27**, 211–221.

Boyd, J. P.: 1998b, *Weakly Nonlocal Solitary Waves and Beyond-All-Orders Asymptotics: Generalized Solitons and Hyperasymptotic Perturbation Theory*, Vol. 442 of *Mathematics and Its Applications*, Kluwer, Amsterdam. 608 pp. Three chapters on spectral methods, solving spectral-discretized nonlinear equations, and special methods for nonlocal solitary waves and other phenomena that radiate to spatial infinity.

Boyd, J. P.: 1998c, High order models for the nonlinear shallow water wave equations on the equatorial beta-plane with application to Kelvin wave frontogenesis, *Dynamics of Atmospheres and Oceans* **28**(2), 69–91.

Boyd, J. P.: 1998d, Two comments on filtering, *J. Comput. Phys.* **143**(1), 283–288. Shows how to apply filters or sum acceleration methods to spectral series so as to preserve the boundary conditions. Also explains why additional boundary conditions are not needed: high order filtering operators can be interpreted as powers of the Legendre or Chebyshev differential operator, which is singular at the boundaries.

Boyd, J. P.: 1998e, Global approximations to the principal real-valued branch of the Lambert W-function, *Appl. Math. Lett.* **11**(6), 27–31.

Boyd, J. P.: 1999a, *Chebyshev and Fourier Spectral Methods*, 2d edn, Dover, Mineola, New York. 665 pp. Heavily revised and updated second edition of Boyd(1989).

Boyd, J. P.: 1999b, A numerical comparison of seven grids for polynomial interpolation on the interval, *Comput. Math. Appl.* Submitted.

Boyd, J. P.: 1999c, The Blasius function in the complex plane, *J. Experimental Math.* In press.

Boyd, J. P.: 2001, *Essays on Chebyshev and Fourier Spectral Methods*. To appear.

Boyd, J. P. and Christidis, Z. D.: 1982, Low wavenumber instability on the equatorial beta-plane, *Geophysical Research Letters* **9**, 769–772.

Boyd, J. P. and Christidis, Z. D.: 1983, Instability on the equatorial beta-plane, *in* J. Nihoul (ed.), *Hydrodynamics of the Equatorial Ocean*, Elsevier, Amsterdam, pp. 339–351.

Boyd, J. P. and Christidis, Z. D.: 1987, The continuous spectrum of equatorial Rossby waves in a shear flow, *Dynamics of Atmospheres and Oceans* **11**, 139–151.

Boyd, J. P. and Flyer, N.: 1999, Compatibility conditions for time-dependent partial differential equations and the the rate of convergence of Chebyshev and Fourier spectral methods, *Comput. Meths. Appl. Mech. Engrg.* In press.

Boyd, J. P. and Haupt, S. E.: 1991, Polycnoidal waves: Spatially periodic generalizations of multiple solitary waves, *in* A. R. Osborne (ed.), *Nonlinear Topics of Ocean Physics: Fermi Summer School, Course LIX*, North-Holland, Amsterdam, pp. 827–856.

Boyd, J. P. and Ma, H.: 1990, Numerical study of elliptical modons by a spectral method, *Journal of Fluid Mechanics* **221**, 597–611.

Boyd, J. P. and Natarov, A.: 1998, A Sturm-Liouville eigenproblem of the Fourth Kind: A critical latitude with equatorial trapping, *Stud. Appl. Math.* **101**, 433–455. Rational TB calculation of logarithmically singular eigenproblems along an infinite interval parallel to the real axis, but shifted from it by a constant.

Boyd, J. P. and Tan, B.: 1998, Vortex crystals and non-existence of non-axisymmetric solitary waves in the Flierl-Petviashvili equation, *Chaos, Solitons and Fractals* **9**, 2007–2021. Double Fourier algorithm for a generalized, two-dimensional quasi-geostrophic equation.

Boyd, J. P. and Tan, B.: 1999, Composite bound states of wide and narrow envelope solitons in the Coupled Schroedinger equations through matched asymptotic expansions, *Nonlinearity.* Submitted.

Brachet, M. E., Meiron, D. I., Orszag, S. A., Nickel, B. G., Morf, R. H. and Frisch, U.: 1983, Small-scale structure of the Taylor-Green vortex, *Journal of Fluid Mechanics* **130**, 411–452. Three-dimensional Fourier computations with exploitation of 64-fold symmetry.

Brandt, A., Fulton, S. R. and Taylor, G. D.: 1985, Improved spectral multigrid methods for periodic elliptic problems, *Journal of Computational Physics* **58**, 96–112.

Braverman, E., Israeli, M., Averbuch, A. and Vozovoi, L.: 1998, A fast 3D Poisson solver of arbitrary order accuracy, *J. Comput. Phys.* **144**(1), 109–136. Fourier spectral method for nonperiodic boundary conditions with subtractions to improve convergence and remedy corner singularities.

Brenier, B., Roux, B. and Bontoux, P.: 1986, Comparaison des méthodes tau-Chebyshev et Galerkin dans l'étude de stabilité des mouvements de convection naturelle. Problème des valeurs propres parasites, *Journal de Méchanique théorique et appliquée* **5**, 95–119.

Bridger, A. F. C. and Stevens, D. E.: 1980, Long atmospheric waves and the polar-plane approximation to the earth's spherical geometry, *Journal of the Atmospheric Sciences* **37**, 534–544.

Briggs, W. L.: 1987, *A Multigrid Tutorial*, SIAM, Philadelphia. 88 pp.

Briggs, W. L. and Henson, V. E.: 1995, *The DFT: An Owner's Manual for the Discrete Fourier Transform*, Society for Industrial and Applied Mathematics, Philadelphia.

Briggs, W. L., Newell, A. C. and Sarie, T.: 1981, The mechanism by which many partial difference equations destabilize, *in* H. Haken (ed.), *Chaos and Order in Nature*, Springer-Verlag, New York, pp. 269–273. Aliasing instability.

Brown, D. L. and Minion, M. L.: 1995, Performance of under-resolved two-dimensional flow simulations, *Journal of Computational Physics* **122**, 165–183.

Brown, J. D., Chu, M. T., Ellison, D. C. and Plemmons, R. J. (eds): 1994, *Proceedings of the Cornelius Lanczos International Centenary Conference*, Society for Industrial and Applied Mathematics, Philadelphia. Collection; many articles on spectral methods.

Browning, G. L., Hack, J. J. and Swarztrauber, P. N.: 1988, A comparison of three numerical methods for solving differential equations on the sphere, *Monthly Weather Review* **117**, 1058–1075. Spherical harmonics versus fourth and sixth order finite differences.

Cai, W., Gottlieb, D. and Harten, A.: 1992a, Cell-averaging Chebyshev methods for hyperbolic problems, *Comput. & Math. Applics.* **24**, 37–49.

Cai, W., Gottlieb, D. and Shu, C.: 1989, Essentially nonoscillatory spectral Fourier methods for shock wave calculation, *Mathematics of Computation* **52**(186), 389–410.

Cai, W., Gottlieb, D. and Shu, C. W.: 1992b, On one-sided filters for spectral Fourier approximation of discontinuous functions, *SIAM Journal of Numerical Analysis* **29**, 905–916.

Cain, A. B., Ferziger, J. H. and Reynolds, W. C.: 1984, Discrete orthogonal function expansions for non-uniform grids using the Fast Fourier transform, *Journal of Computational Physics* **56**, 272–286. Use Fourier series with a mapping which is equivalent to rational Chebyshev.

Calahan, D. A.: 1967, Numerical solution of linear systems with widely separated time constants, *Proceedings IEEE* **55**, 2016–2017. Not spectral; higher order Padé time marching scheme.

Canuto, C. and Funaro, D.: 1988, The Schwarz algorithm for spectral methods, *SIAM Journal of Numerical Analysis* **25**, 24–40.

Canuto, C. and Quarteroni, A.: 1985, Preconditioned minimal residual methods for Chebyshev spectral calculations, *Journal of Computational Physics* **60**, 315–337.

Canuto, C., Hussaini, M. Y., Quarteroni, A. and Zang, T. A.: 1988, *Spectral Methods for Fluid Dynamics*, Springer-Verlag, New York. Classic text, 556 pp. Very comprehensive and readable, now in paperback.

Carcione, J. M.: 1994, Boundary conditions for wave propagation problems, *in* C. Bernardi and Y. Maday (eds), *Analysis, Algorithms and Applications of Spectral and High Order Methods for Partial Differential Equations*, Selected Papers from the International Conference on Spectral and High Order Methods (ICOSAHOM '92), Le Corum, Montpellier, France, 22-26 June 1992, North-Holland, Amsterdam, pp. 457–468. Also in Finite Elements in Analysis and Design, vo. 16, pp. 317-327. Compares standard Chebyshev grid with the stretched Kosloff/Tal-Ezer grid.

Carcione, J. M.: 1996, A 2-D Chebyshev differential operator for the elastic wave equation, *Computer Methods in Applied Mechanics and Engineering* **130**, 33–45. Adaptive grid mapping using the Kosloff/Tal-Ezer map (for longer time step), the Augenbaum map (to resolve narrow features) and different maps on different subdomains for maximum flexibility.

Carpenter, M. H.: 1996, Spectral methods on arbitrary grids, *Journal of Computational Physics* **129**(1), 74–86. Differentiation is performed using one grid of points while the equation is collocated on a different grid. Generalization of Don and Gottlieb(1994).

Carpenter, R. L., Droegemeier, K. K., Woodward, P. R. and Hane, C. E.: 1990, Application of the piecewise parabolic method (PPM) to meteorological modeling, *Monthly Weather Review* **118**, 586–612. Not spectral.

Carrier, G. F. and Pearson, C. E.: 1968, *Ordinary Differential Equations*, Blaisdell, Waltham, MA. 229 pp; not spectral.

Carse, G. A. and Urquhart, J.: 1914, Harmonic analysis, *in* E. M. Horsburgh (ed.), *Modern Instruments and Methods of Calculation*, G. Bell and Sons, in cooperation with the Royal Society of Edinburgh, London. 300 pp. Reviews "Runge grouping", which is a variant of the FFT that was widely used in the early 20th century and requires only real arithmetic. This algorithm was lost in mid-century until rediscovered, in complex form, by Cooley and Tukey. The article also describes the Runge-FFT computing forms devised by E. T. Whittaker [today, we would call them "spreadsheets"] and the mechanical harmonic analyzer of Michaelson and Stratton (1898) which empirically discovered the Gibbs' phenomenon.

Chan, T. and Kerkhoven, T.: 1985, Fourier methods with extended stability intervals for the Korteweg-deVries equation, *SIAM Journal of Numerical Analysis* **22**, 441–454.

Chan, T. F.: 1984, Newton-like pseudo-arclength methods for computing simple turning points, *SIAM Journal of Scientific and Statistical Computing* **5**, 135–148. Not spectral, but useful for tracking branches of the solutions to nonlinear equations as a parameter is varied.

Chaouche, A. M.: 1990, A collocation method basedon an influence matrix technique for axisymmetric flows in an annulus, *Rech. Aérosp.* **1990**-**5**, 1–13.

Chaouche, A., Randriamampianina, A. and Bontoux, P.: 1990, A collocation method based on an influence matrix technique for axisymmetric flows in an annulus, *Computer Methods in Applied Mechanics and Engineering* **80**, 237–244.

Chapman, S. and Lindzen, R. S.: 1970, *Atmospheric Tides*, D. Reidel, Dordrecht, Holland. 200 pp., computes Hough functions by solving eigenvalue problem for Laplace's tidal equations by spherical harmonic expanions.

Chen, H. B.: 1993a, On the instability of a full non-parallel flow — Kovasznay flow, *International Journal for Numerical Methods in Fluids* **17**, 731–754. Eigenvalue study using rational Chebyshev functions TL_n for the semi-infinite interval.

Chen, S., Doolen, G. D., Kraichnan, R. H. and She, Z.-S.: 1993, On statistical correlation between velocity increments and locally averaged dissipation in homogeneous turbulence, *Physics of Fluids A* **5**, 458–463.

Chen, X.-S.: 1993b, The aliased and dealiased spectral models of the shallow-water equations, *Monthly Weather Review* **121**, 834–852. Solves the KdV equation and also the spherical shallow water wave equations for both very smooth initial conditions (Rossby-Haurwitz waves) and realistic conditions (observational data from FGGE). In all cases, the aliased models are no better than aliased models with the same number of points after filtering (the dealiased code uses $(3/2)N$ points to compute N modes); the aliased models are always considerably worse than dealiased models when compared on the basis of the same number of collocation points (the dealiased code has a smaller number of modes than the aliased code after the dealiasing filtering is applied). One of the dealiased methods is novel in that it uses the Walsh Hadamard transform instead of the usual FFT. The WT transform method is a little cheaper than the FFT for the same accuracy, but all of the dealiased codes are 1. 7 to 2 times more expensive than the aliasing codes with the same number of grid points. but unfortunately no more accurate.

Christov, C. I.: 1982, A complete orthonormal system in $L^2(-\infty, \infty)$ space, *SIAM Journal of Applied Mathematics* **42**, 1337–1344.

Christov, C. I. and Bekyarov, K. L.: 1990, A Fourier-series method for solving soliton problems, *SIAM Journal of Scientific and Statistical Computing* **11**, 631–647. Rational functions.

Chu, M. T.: 1988, On the continuous realization of iterative processes, *SIAM Review* **30**, 375–387. Differential equations in pseudotime as models for Newton's and other iterations.

Cividini, A. and Zampieri, E.: 1997, Nonlinear stress analysis problems by spectral collocation methods, *Comput. Methods Appl. Mech. Engrg.* **145**, 185–201.

Clenshaw, C. W.: 1957, The numerical solution of linear differential equations in Chebyshev series, *Proceedings of the Cambridge Philosophical Society* **53**, 134–149.

Clenshaw, C. W. and Curtis, A. R.: 1960, A method for numerical integration on an automatic computer, *Numerische Mathematik* **2**, 197–205. Chebyshev polynomial quadrature scheme.

Clercx, H. J. H.: 1997, A spectral solver for the Navier-Stokes equations in the velocity-vorticity formulation for flows with two nonperiodic directions, *J. Comput. Phys.* **137**, 186–211.

Cloot, A.: 1991, Equidistributing mapping and spectral method for the computation on unbounded domains, *Applied Mathematics Letters* **4**, 23–27.

Cloot, A. and Weideman, J. A. C.: 1990, Spectral methods and mappings for evolution equations on the infinite line, *Computer Methods in Applied Mechanics and Engineering* **80**, 467–481.

Cloot, A. and Weideman, J. A. C.: 1992, An adaptive algorithm for spectral computations on unbounded domains, *Journal of Computational Physics* **102**, 398–406.

Cloot, A., Herbst, B. M. and Weideman, J. A. C.: 1990, A numerical study of the cubic-quintic Schrodinger equation, *Journal of Computational Physics* **86**, 127–146.

Concus, P. and Golub, G. H.: 1973, Use of fast direct methods for the efficient numerical solution of nonseparable elliptic equations, *SIAM Journal of Numerical Analysis*. Not spectral, but shows how a fast direct method, either spectral or otherwise, can be used to efficiently solve nonseparable elliptic equations.

Cooley, J. W. and Tukey, J. W.: 1965, An algorithm for the machine calculation of complex Fourier series, *Mathematics of Computation* **19**, 297–301.

Corral, R. and Jiménez, J.: 1995, Fourier/Chebyshev methods for the incompressible Navier-Stokes equations in infinite domains, *J. Comput. Phys.* **121**, 261.

Côté, J., Gravel, S. and Staniforth, A.: 1990, Improving variable-resolution finite-element semi-Lagrangian integration schemes by pseudostaggering, *Monthly Weather Review* **118**, 2718–2731.

Côté, J., Gravel, S. and Staniforth, A.: 1995, A generalized family of schemes that eliminates the spurious resonant response of semi-Lagrangian schemes to orographic forcing, *Monthly Weather Review* **123**, 3605.

Coulson, C. A.: 1961, *Valence*, Oxford University Press, New York. 250 pp.

Courtier, P. and Geleyn, J.-F.: 1988, A global numerical weather prediction model with variable resolution: Application to the shallow-water equations, *Quarterly Journal of Royal Meteorological Society* **114**, 1321–1346.

Courtier, P. and Naughton, M.: 1994, A polar problem in the reduced Gaussian grid, *Quart. J. Roy. Met. Soc.* **120**, 1389–1407.

Courtier, P., Freydier, C., Geleyn, J. F., Rabier, F. and Rochas, M.: 1991, The Arpege project at Météo-France., *Proceedings of Numerical Methods in Atmospheric Models*, European Center for Medium Range Forecasting, European Center for Medium Range Forecasting, Shinfield Park, Reading, United Kingdom, pp. 193–231.

Coutsias, E. A., Hagstrom, T. and Torres, D.: 1996, An efficient spectral method for ordinary differential equations with rational function coefficients, *Mathematics of Computation* **65**, 611–635. Obtained banded matrix representations of one-dimensional differential operators for all standard polynomial basis sets, assuming that the coefficients of the operators are restricted to polynomials or rational functions. These banded matrices can be inverted in $O(N)$ operations.

Couzy, W. and Deville, M.: 1994, Iterative solution technique for spectral-element pressure operators at high Reynolds number, *in* S. Wagner et al. (eds), *Proceedings of the Second European Computational Fluid Dynamics Conference*, Stuttgart, Germany, pp. 613–618.

Craik, A. D. D.: 1985, *Wave Interactions and Fluid Flows*, Cambridge University Press, New York. Not spectral, but a good description of weakly nonlinear waves and resonant triad and four-wave interactions.

Curchitser, E. N., Iskandarani, M. and Haidvogel, D. B.: 1998, A spectral element solution of the shallow-water equations on multiprocessor computers, *J. Atmos. Oceanic Technology* **15**(2), 510–521.

da Silva, M. R.: 1987, A quick survey of recent developments and applications of the tau-method, *in* E. L. Ortiz (ed.), *Numerical Approximations of P. D. E., Part III*, North-Holland, Amsterdam, pp. 297–308.

Daley, R.: 1980, The development of efficient time integration schemes using model normal modes, *Monthly Weather Review* **108**, 100–110.

Daley, R.: 1981, Normal mode initialization, *Reviews of Geophysics and Space Physics* **19**(3), 450–468. Review.

Daley, R.: 1991, *Atmospheric Data Analysis*, Cambridge University Press, New York.

Daley, R. and Bourassu, Y.: 1978, Rhomboidal versus triangular spherical harmonic truncation: Some verification statistics, *Atmosphere-Ocean* **16**, 187–196.

Davis, P. J.: 1975, *Interpolation and Approximation*, Dover Publications, New York. 200 pp.

Davis, P. J. and Rabinowitz, P.: 1984, *Methods of Numerical Integration*, 3rd edn, Academic Press, Boston. 612pp.

Dawkins, P. T., Dunbar, S. R. and Douglass, R. W.: 1998, The origin and nature of spurious eigenvalues in the spectral tau method, *J. Comput. Phys.* **147**(2), 441–462. Theoretical paper proving the existence of two spurious eigenvalues for the exemplary problem $u_{xxxx} = \lambda u_{xx}$ which are larger than N^4. The cure and numerical experiments are also described.

de Veronico, M. C., Funaro, D. and Reali, G. C.: 1994, A novel numerical technique to investigate nonlinear guided waves: approximation of the Nonlinear Schroedinger equation by nonperiodic pseudospectral methods, *Numerical Methods for Partial Differential Equations* **10**(6), 667–675. Domain decomposition with Legendre polynomials on interior domains and Laguerre polynomials on the exterior elements, which extend to infinity.

de Vries, R. W. and Zandbergen, P. J.: 1989, The numerical solution of the biharmonic equation, using a spectral multigrid method, *in* W. F. Ballhaus and M. Y. Hussaini (eds), *Advances in Fluid Dynamics*, Springer-Verlag, New York, pp. 25–35.

Deane, A. E., Kevrekidis, I. G., Karniadakis, G. E. and Orszag, S. A.: 1991, Low-dimensional modeling for complex geometry flows: Application to grooved channels and circular cylinders, *Physics of Fluids A* **3**, 2337–2354. Spectral elements; also derivation and application of a four-mode model using "empirical eigenfunctions" as the basis (also known as "proper orthogonal decomposition".

Debussche, A., Dubois, T. and Temam, R.: 1995, The Nonlinear Galerkin method: A multiscale method applied to the simulation of homogeneous turbulent flows, *Theoretical and Computational Fluid Dynamics* **7**, 279–315.

Decker, D. W. and Keller, H. B.: 1980, Path following near bifurcation, *Communications in Pure and Applied Mathematics* **34**, 149–175.

Delves, L. M.: 1976, Expansion methods, *in* G. Hall and J. M. Watts (eds), *Modern Numerical Methods for O. D. E.s*, Clarendon Press, Oxford University Press, Oxford, pp. 269–281. REVIEW.

Delves, L. M.: 1977a, A fast method for the solution of Fredholm integral equations, *Journal of the Institute for Mathematics and its Applications* **20**, 173–184.

Delves, L. M.: 1977b, On the solution of the linear equation arising from Galerkin methods, *Journal of the Institute for Mathematics and its Applications* **20**, 163–171.

Delves, L. M.: 1977c, A linear equation solver for Galerkin and least squares methods, *J. Comp.* **20**, 371–374.

Delves, L. M., Abd-Elal, L. F. and Hendry, J. A.: 1979, A fast Galerkin algorithm for the solution of Fredholm integral equations, *Journal of the Institute for Mathematics and its Applications* **23**, 139–166.

Delves, L. M., Abd-Elal, L. F. and Hendry, J. A.: 1981a, A set of modules for the solution of integral equations, *Comp. J.* **24**, 184–190.

Delves, L. M. and Hall, C. A.: 1979, An implicit matching procedure for global element calculations, *Journal of the Institute for Mathematics and its Applications* **23**, 223–234.

Delves, L. M. and Mead, K. O.: 1971, On the convergence rates of variational methods. I. Asymptotically diagonal systems, *Mathematics of Computation* **25**, 699–716. Theory only.

Delves, L. M. and Phillips, C.: 1980, A fast implementation of the global element method, *Journal of the Institute for Mathematics and its Applications* **25**, 177–197.

Delves, L. M., McKerrell, A. and Henry, J. A.: 1981b, A note on Chebyshev methods for the solution of partial differential equations, *Journal of Computational Physics* **41**, 444–452.

Delves, L. M., McKerrell, A. and Peters, S. A.: 1986, Performance of GEM2 on the ELLPACK problem population, *International Journal for Numerical Methods in Engineering* **23**, 229–238.

Delves, L. N. and Freeman, T. N.: 1981, *Analysis of Global Expansion Methods: Weakly Asymptotically Diagonal Systems*, Academic Press, New York. 275 pp. Mostly theory with only a handful of elementary examples, but the preconditioned "Delves-Freeman" iteration is very interesting.

Demaret, P. and Deville, M. O.: 1989, Chebyshev pseudospectral solution of the Stokes equation using finite element preconditioning, *Journal of Computational Physics* **83**, 463–484.

Demaret, P. and Deville, M. O.: 1991, Chebyshev collocations solutions of the Navier-Stokes equations using multi-domain decomposition and finite element preconditioning, *Journal of Computational Physics* **95**, 359–386.

Demaret, P., Deville, M. O. and Schneidesch, C.: 1989, Thermal convection solutions by Chebyshev pseudospectral multi-domain decomposition and finite element preconditioning, *Applied Numerical Mathematics* **6**, 107–121. Nonlinear steady flows through nested iterations: outer Newton/inner Richardson.

Déqué Piedelievre, M.: 1995, High resolution climate model over Europe, *Climate Dynamics* **11**, 321–339.

Dettori, L., Gottlieb, D. and Témam, R.: 1995, Nonlinear Galerkin method: the two-level Fourier collocation case, *Journal of Scientific Computing* **10**, 371–.

Deville, M.: 1990, Chebyshev collocation solutions of flow problems, *in* C. Canuto and A. Quarteroni (eds), *Spectral and High Order Methods for Partial Differential Equations: Proceedings of the ICOSAHOM '89 Conference in Como, Italy*, North-Holland/Elsevier, Amsterdam, pp. 27–38. Also in Comput. Meths. Appl. Mech. Engrg., vol. 80.

Deville, M. and Labrosse, G.: 1982, An algorithm for the evaluation of multi-dimensional (direct and inverse) discrete Chebyshev transform, *Journal of Computational and Applied Mathematics* **8**, 293–304.

Deville, M. and Mund, E.: 1984, On a mixed one step/Chebyshev pseudospectral technique for the integration of parabolic problems using finite element preconditioning, *in* C. Brezinski, A. Draux, A. P. Magnus, P. Maroni and A. Ronveaux (eds), *Polynomes Orthogonaux et Applications: Proceedings of the Laguerre Symposium at Bar-le-Duc*, number 1171 in *Lecture Notes in Mathematics*, Springer-Verlag, New York, pp. 399–407. This article is in English; employs unusual implicit time-marching which is exact (instead of second order) for time integration of the two slowest-decaying diffusion eigenmodes.

Deville, M. and Mund, E.: 1985, Chebyshev pseudospectral solution of second-order elliptic equations with finite element pre-conditioning, *Journal of Computational Physics* **60**, 517–533.

Deville, M. and Mund, E.: 1991, Finite element preconditioning of collocation schemes for advection-diffusion equations, *in* R. Beauwens and P. de Groen (eds), *Proceedings of the IMACS International Symposium on Iterative Methods in Linear Algebra*, IMACS, North-Holland, Amsterdam, pp. 181–190.

Deville, M., Haldenwang, P. and Labrosse, G.: 1981, Comparison of time integration (finite difference and spectral)) for the nonlinear Burgers' equation, *in* H. Viviond (ed.), *Proceedings of the 4th GAMMConference on Nuemrical Methods in Fluid Mechanics*, Vieweg, Braunschweig.

Deville, M., Kleiser, L. and Montigny-Rannou, F.: 1984, Pressure and time treatment for Chebyshev spectral solution of a Stokes problem, *Internat. J. Numer. Meth. Fluids* **4**, 1149–1163. Backward Euler is raised by Richardson extrapolation to a second order time-marching. Four different schemes including penalty method, splitting, influence matrix and Morchoisne's space-time pseudospectral scheme. Good discussion of compatibility conditions on the initial condition.

Deville, M. O. and Mund, E. H.: 1990, Finite-element preconditioning for pseudospectral solutions of elliptic problems, *SIAM Journal of Scientific and Statistical Computing* **12**, 311–342.

Deville, M. O. and Mund, E. H.: 1992, Fourier analysis of finite element preconditioned collocation schemes, *SIAM Journal of Scientific and Statistical Computing* **13**(2), 596–610.

Deville, M. O., Mund, E. H. and Van Kemenade, V.: 1994, Preconditioned Chebyshev collocation methods and triangular finite elements, *in* C. Bernardi and Y. Maday (eds),

Analysis, Algorithms and Applications of Spectral and High Order Methods for Partial Differential Equations, Selected Papers from the International Conference on Spectral and High Order Methods (ICOSAHOM '92), Le Corum, Montpellier, France, 22-26 June 1992, North-Holland, Amsterdam, pp. 193–200.

Devulder, C. and Marion, M.: 1992, A class of numerical algorithms for large time integration: the nonlinear Galerkin methods, *SIAM Journal of Numerical Analysis* **29**(2), 462–483. Many theorems.

Dickinson, B. A.: 1933, Energy levels of H_2^+ molecular ion, *J. Chem. Phys.* **1**, 317. Illustration of the Rayleigh variational principle in quantum chemistry;.

Dimitropoulus, C. and Beris, A. N.: 1997, An efficient and robust spectral solver for non-separable elliptic equations, *Journal of Computational Physics* **133**, 186–191. Biconjugate gradient outer iteration with a Concus-Golub inner iteration in which a separable Helmholtz equation is solved at each iteration.

Dimitropoulus, C. and Beris, A. N.: 1998, Efficient pseudospectral flow simulations in moderately complex geometries, *Journal of Computational Physics* **144**(2), 517–549. Pseudoconformal orthogonal curvilinear coordinates in two dimensions, Fourier in one coordinate and Chebyshev in the other.

Don, W. S. and Gottlieb, D.: 1990, Spectral simulation of unsteady flow past a cylinder, *in* C. Canuto and A. Quarteroni (eds), *Spectral and High Order Methods for Partial Differential Equations: Proceedings of the ICOSAHOM '89 Conference in Como, Italy*, North-Holland/Elsevier, Amsterdam, pp. 39–58. Also in Comput. Meths. Appl. Mech. Engrg., vol. 80, with the same page numbers.

Don, W. S. and Gottlieb, D.: 1994, The Chebyshev-Legendre method: implementing Legendre methods on Chebyshev points, *SIAM Journal of Numerical Analysis* **31**, 1519–1534.

Don, W. S. and Solomonoff, A.: 1995, Accuracy and speed in computing the Chebyshev collocation derivative, *SIAM J. Sci. Comput.* **16**, 1253–1268.

Don, W. S. and Solomonoff, A.: 1997, Accuracy enhancement for higher derivatives using Chebyshev collocation and a mapping technique, *SIAM Journal of Scientific Computing* **18**(4), 1040–1055. The Kosloff/Tal-Ezer mapping is used to reduce roundoff error and allow a larger timestep.

Dongarra, J. J., Straughan, B. and Walker, D. W.: 1996, Chebyshev tau-QZ algorithm methods for calculating spectra of hydrodynamic stability problems, *Applied Numerical Mathematics* **22**, 399–434.

Doron, E., Hollingsworth, A., Hoskins, B. J. and Simmons, A. J.: 1974, A comparison of grid-point and spectral methods in a meteorological problem, *Quarterly Journal of the Royal Meteorological Society* **100**, 371–383.

Douglas, J. and Russell, T. F.: 1982, Numerical methods for convection-dominated diffusion problems based on combining the method of characteristics with finite element or finite difference procedures, *SIAM Journal of Numerical Analysis* **19**, 871–885. Not spectral; invention of a semi-Lagrangian scheme.

Drake, J., Foster, I., Michalakes, J., Toonen, B. and Worley, P.: 1995, Design and performance of a scalable parallel community climate model, *Parallel Comput.* **21**(10), 1571–1591. Parallel version, PCCM2, of the CCM2 spherical harmonics/vertical finite difference climate model. Performance on the IBM SP2 and Intel Paragon.

Driscoll, J., Healy, Jr., D. M. and Rockmore, D.: 1997, Fast discrete polynomial transform with applications to data analysis on distance transitive graphs, *SIAM J. Comput.* **26**, 1066–1099. $O(N \log_2(N))$ transform which works for any set of orthogonal polynomials; more efficient than Orszag(1986).

Driscoll, T. A. and Fornberg, B.: 1998, A block pseudospectral method for Maxwell's equations. I. One-dimensional case, *Journal of Computational Physics* **140**(1), 47–65. Employ a domain decomposition scheme in which fictitious points beyond the domain walls are used, in a sort of grid overlapping scheme, to allow a much more uniform separation between grid points than in a standard pseudospectral algorithm. This allows a relatively long time step at the cost of much increased domain-to-domain communication compared to the standard domain decomposition method. They generalize the scheme so that it works well even when there are discontinuous changes in material properties at domain walls.

Dubois, T., Jauberteau, F. and Témam, R.: 1990, The nonlinear Galerkin method for the two and three dimensional Navier-Stokes equations, *in* K. W. Morton (ed.), *Proceedings of the Twelfth International Conference on Numerical Methods in Fluid Dynamics*, Springer-Verlag, New York, pp. 117–120.

Dubois, T., Jauberteau, F. and Témam, R.: 1998, Incremental unknowns, multilevel methods and the numerical simulation of turbulence, *Comput. Meth. Appl. M.* **159**, 123–189.

Durran, D. R.: 1991, The third-order Adams-Bashforth method: an attractive alternative to leapfrog time-differencing, *Monthly Weather Review* **119**, 702–720.

Dutt, A. and Rokhlin, V.: 1993, Fast Fourier Transforms for nonequispaced data, *SIAM J Comput.* **14**, 1368–1393.

Dutt, A. and Rokhlin, V.: 1995, Fast Fourier Transforms for nonequispaced data, II, *Applied and Computational Harmonic Analysis* **2**, 85–110.

D'yakonov, E. G.: 1961, An iteration method for solving systems of finite difference equations, *Doklady Akademiia Nauk SSSR* **138**, 522–525. Not spectral; iteration preconditioned by separable PDE.

Dym, H. and McKean, H. P.: 1972, *Fourier Series and Integrals*, Academic Press, New York. 129 pp.

Eggert, N., Jarratt, M. and Lund, J.: 1987, Sinc function computation of Sturm-Liouville problems, *Journal of Computational Physics* **69**, 209–229.

Ehrenstein, U. and Peyret, R.: 1989, A Chebyshev collocation method for the Navier-Stokes equations with application to double-diffusive convection, *International Journal for Numerical Methods in Fluids* **9**, 427–452. Semi-implicit time integration with influence matrix method.

Eisen, H. and Heinrichs, W.: 1992, A new method of stabilization for singular perturbation problems with spectral methods, *SIAM Journal of Numerical Analysis* **29**, 107–122. Shows that basis functions which vanish at boundaries are much better conditioned than the Chebyshev polynomials from which these basis functions are formed.

Eisen, H., Heinrichs, W. and Witch, K.: 1991, Spectral collocation methods and polar coordinate singularities, *Journal of Computational Physics* **96**, 241–257.

Eisenstat, S. C., Elman, H. C. and Schultz, M. H.: 1983, Variational iterative methods for nonsymmetric systems of linear equations, *SIAM J. Numer. Anal.* **20**, 345–357.

El-Daou, M. K. and Ortiz, E. L.: 1993, Error analysis of the tau method: dependence of the error on the degree and on the length of the interval of approximation, *Computers Math. Applic.* **25**(7), 33–45.

El-Daou, M. K. and Ortiz, E. L.: 1994a, A recursive formulation of collocation in terms of canonical polynomials, *Computing* **52**, 177–202.

El-Daou, M. K. and Ortiz, E. L.: 1994b, The weighting subspaces of collocation and the Tau method, *in* J. D. Brown, M. T. Chu, D. C. Ellison and R. J. Plemmons (eds), *Proceedings of the Cornelius Lanczos International Centenary Conference*, Society for Industrial and Applied Mathematics, Society for Industrial and Applied Mathematics, Philadelphia, PA.

El-Daou, M. K. and Ortiz, E. L.: 1997, The uniform convergence of the Tau method for singularly perturbed problems, *Applied Mathematics Letters* **10**(2), 91–94. Existence and stability of the algorithm is proved, independent of the perturbation parameter ϵ.

El-Daou, M. K., Ortiz, E. L. and Samara, H.: 1993, A unified approach to the tau method and Chebyshev series expansion techniques, *Computers Math. Applic.* **25**(3), 73–82.

Eliassen, E. and Machenhauer, B.: 1974, On spectral representation of the vertical variation of the meteorological fields in numerical integration of a primitive equation model, *GARP WGNE Report 7*, World Meteorological Organization, Geneva, Switzerland. Legendre polynomials in the vertical.

Elliott, D. and Stenger, F.: 1984, Sinc method of solution of singular integral equations, *in* A. Gerasoulis and R. Vichnevetsky (eds), *Numerical Solution of Singular Integral Equations*, IMACS.

Ellsaesser, H. W.: 1966, Evaluation of spectral versus grid point methods of hemispheric numerical weather prediction, *J. Appl. Meteor.* **5**, 246–262. Fig. 12 is a good illustration of late onset of spectral blocking due to violation of the CFL criterion after the advecting flow has intensified from its initial maximum.

Engelmann, F., Feix, M., Minardi, E. and Oxenius, J.: 1963, Nonlinear effects from vlasov's equation, *The Physics of Fluids* **6**(2), 266–275. Low order Hermite series ($N = 2, 3$) for velocity coordinate; such low truncations are found to suppress instabilities that occur in some parameter regions for the full equations.

Erlebacher, G., Zang, T. A. and Hussaini, M. Y.: 1987, Spectral multigrid methods for the numerical simulation of turbulence, *in* S. McCormick and K. Stuben (eds), *Multigrid Methods*, Marcel Dekker, New York, pp. 177–194.

Errico, R. M.: 1984, The dynamic balance of a general circulation model, *Monthly Weather Review* **112**, 2439–2454. Not spectral, but evaluation of the usefulness of slow manifold concept in a complicated model.

Errico, R. M.: 1989, The degree of Machenauer balance in a climate model, *Monthly Weather Review* **112**, 2723–2733. Test of slow manifold ideas of Machenauer's NG(1) approximation in a global hydrodynamics-with-physics model.

Falqués, A. and Iranzo, V.: 1992, Edge waves on a longshore shear flow, *Physics of Fluids* pp. 2169–2190. Rational Chebyshev and Laguerre on semi-infinite domain.

Finlayson, B. A.: 1973, *The Method of Weighted Residuals and Variational Principles*, Academic, New York. 412 pp. Many good examples of low order pseudospectral methods, dubbed "orthogonal collocation" here, and mostly drawn from chemical engineering and fluid dynamics. Mostly Legendre and Gegenbauer polynomials rather than Chebyshev and no mention of the FFT.

Fischer, P. F.: 1990, Analysis and application of a parallel spectral element method for the solution of the Navier-Stokes equations, *Computer Methods in Applied Mechanics and Engineering* **80**(1–3), 483–491.

Fischer, P. F.: 1994a, Domain decomposition methods for large scale parallel Navier-Stokes calculations, *in* A. Quarteroni (ed.), *Proceedings of the Sixth International Conference on Domain Decomposition Methods for Partial Differential Equations, Como, Italy*, AMS, Providence.

Fischer, P. F.: 1994b, Parallel domain decomposition for incompressible fluid dynamics, *Contemp. Math.* **157**, 313.

Fischer, P. F.: 1997, An overlapping Schwarz method for spectral element solution of the incompressible Navier-Stokes equations, *J. Comput. Phys.* **133**, 84–101.

Fischer, P. F.: 1998, Projection techniques for iterative solution of ax=b with successive right-hand sides, *Comput. Meth. Appl. M.* **163**(1–4), 193–204. Not spectral, but useful for semi-implicit time marching.

Fischer, P. F. and Gottlieb, D.: 1997, On the optimal number of subdomains for hyperbolic problems on parallel computers, *International Journal of Supercomputing and Appl. High Performance Computing* **11**, 65–76. Spectral elements.

Fischer, P. F. and Patera, A. T.: 1989, Parallel spectral element methods for the incompressible Navier-Stokes equations, *in* J. H. Kane and A. D. Carlson (eds), *Solution of Super Large Problems in Computational Mechanics*, Plenum, New York.

Fischer, P. F. and Patera, A. T.: 1991, Parallel spectral element solution of the Stokes problem, *Journal of Computational Physics* **92**(2), 380–421.

Fischer, P. F. and Patera, A. T.: 1992, Parallel spectral element solutions of eddy-promoter channel flow, *Proceedings of the European Research Community on Flow Turbulence and Combustion Workshop, Laussane, Switzerland*, Cambridge University Press, Cambridge.

Fischer, P. F. and Patera, A. T.: 1994, Parallel simulation of viscous incompressible flows, *Annual Reviews of Fluid Mechanics* **26**, 483–527. REVIEW.

Fischer, P. F. and Rønquist, E. M.: 1994, Spectral element methods for large scale parallel Navier-Stokes calculations, *in* C. Bernardi and Y. Maday (eds), *Analysis, Algorithms and Applications of Spectral and High Order Methods for Partial Differential Equations*, Selected Papers from the International Conference on Spectral and High Order Methods (ICOSAHOM '92), Le Corum, Montpellier, France, 22-26 June 1992, North-Holland, Amsterdam, pp. 69–76. Also in Comput. Methods. Appl. Mech. Engrg., vol. 116.

Fischer, P. F., Ho, L.-W., Karniadakis, G. E., Rønquist, E. M. and Patera, A. T.: 1988a, Recent advances in parallel spectral element simulation of unsteady incompressible flows, *Computers and Structures* **30**, 217–231.

Fischer, P. F., Rønquist, E. M. and Patera, A. T.: 1989, Parallel spectral element methods for viscous flow, *in* G. Carey (ed.), *Parallel Supercomputing: Methods, Algorithms and Applications*, Wiley, New York, pp. 223–238.

Fischer, P., Rønquist, E. M., Dewey, D. and Patera, A. T.: 1988b, Spectral element methods: Algorithms and architectures, *in* R. Glowinski, G. Golub, G. Meurant and J. Periaux (eds), *Proceedings of the First International Conference on Domain Decomposition Methods for Partial Differential Equations*, SIAM, SIAM, Philadelphia, pp. 173–197.

Fjørtoft, R.: 1952, On a numerical method of integrating the barotropic vorticity equation, *Tellus* **4**, 179–194. Not spectral; early use of Lagrangian coordinates in numerical meteorology.

Fjørtoft, R.: 1955, On a numerical method of integrating the barotropic vorticity equation, *Tellus* **7**, 462–480. Not spectral; early use of Lagrangian coordinates in numerical meteorology.

Flå, T.: 1992, A numerical energy conserving method for the DNLS equation, *Journal of Computational Physics* **101**, 71–79. Fourier pseudospectral scheme for the Derivative-Nonlinear Schroedinger equation.

Flatau, P., Boyd, J. P. and Cotton, W. R.: 1987, Symbolic algebra in applied mathematics and geophysical fluid dynamics — REDUCE examples, *Technical report*, Colorado State University, Department of Atmospheric Science, Fort Collins, CO 80523. Some examples of spectral methods in REDUCE.

Flyer, N.: 1998, Asymptotic upper bounds for the coefficients in the Chebyshev series expansion for a general order integral of a function, *Math. Comput.* **67**(224), 1601–1616.

Foias, C., Jolly, M. S., Kevrekidis, I. G. and Titi, E. S.: 1991, Dissipativity of numerical schemes, *Nonlinearity* **4**, 591–613. Blow-up of various schemes including Nonlinear Galerkin.

Foias, C., Jolly, M. S., Kevrekidis, I. G. and Titi, E. S.: 1994, On some dissipative fully discrete nonlinear Galerkin schemes for the Kuramoto-Sivashinsky equation, *Physics Letters A* **186**, 87–96.

Foias, C., Manley, O. and Témam, R.: 1988, Modelling of the interaction of small and large eddies in two dimensional turbulent flows, *Rairo-Math. Mod. Numer. Anal.* **22**(1), 93–118. First paper on approximate inertial manifolds, which led to rediscovery of Nonlinear Galerkin method.

Forbes, J. M. and Garrett, H. B.: 1976, Solar diurnal tide in the thermosphere, *Journal of the Atmospheric Sciences* **33**, 2226–2241.

Fornberg, B.: 1975, On a Fourier method for the integration of hyperbolic equations, *SIAM Journal of Numerical Analysis* **12**, 509–528.

Fornberg, B.: 1977, A numerical study of 2-d turbulence, *Journal of Computer Physics* **25**, 1–31.

Fornberg, B.: 1978, Pseudospectral calculations on 2-d turbulence and nonlinear waves, *SIAM-AMS Proceedings* **11**, 1–18.

Fornberg, B.: 1987, The pseudospectral method: Comparisons with finite differences for the elastic wave equation, *Geophysics* **52**, 483–501.

Fornberg, B.: 1988a, The pseudospectral method: accurate representation of interfaces in elastic wave calculations, *Geophysics* **53**, 625–637.

Fornberg, B.: 1988b, Generation of finite difference formulas on arbitrarily spaced grids, *Mathematics of Computation* **51**, 699–706. Simple recursion to compute finite difference weights. This is helpful for finite difference preconditioning of spectral methods; the finite difference problem must be solved on the same unevenly spaced grid as used by the Chebyshev or Legendre pseudospectral scheme.

Fornberg, B.: 1990a, High order finite differences and the pseudospectral method on staggered grids, *SIAM Journal of Numerical Analysis* **27**, 904–918.

Fornberg, B.: 1990b, An improved pseudospectral method for initial boundary value problems, *Journal of Computational Physics* **91**, 381–397. Uses additional boundary conditions, derived from the differential equation, to greatly reduce the largest eigenvalues of Chebyshev differentiation matrices, greatly reducing the "stiffness" of time-dependent problems and allowing a longer time step.

Fornberg, B.: 1992, Fast generation of weights in finite difference formulas, *in* G. D. Byrne and W. E. Schiesser (eds), *Recent Developments in Numerical Methods and Software for QDEs/DAEs/PDEs*, World Scientific, Singapore, pp. 97–123.

Fornberg, B.: 1995, A pseudospectral approach for polar and spherical geometries, *SIAM Journal of Scientific Computing* **16**, 1071–1081. Double Fourier series on latitude and longitude grid with strong filtering of large zonal wavenumbers near the poles to avoid CFL instability.

Fornberg, B.: 1996, *A Practical Guide to Pseudospectral Methods*, Cambridge University Press, New York.

Fornberg, B.: 1998, Calculation of weights in finite difference formulas, *SIAM Rev.* **40**(3), 685–691. Arbitrary order finite differences, which in the limit of order equal to the number of grid points gives a pseudospectral approximation.

Fornberg, B. and Merrill, D.: 1997, Comparison of finite difference and pseudospectral methods for convective flow over a sphere, *Geophys. Res. Lett.* **24**, 3245–3248. Second and fourth order finite differences and double Fourier series, all on a latitude-longitude grid with strong high latitude filtering of large zonal wavenumbers, are compared with spherical harmonics for a problem whose solution is very smooth. Equal accuracy for both spectral methods, but the double Fourier algorithm is faster.

Fornberg, B. and Sloan, D.: 1994, A review of pseudospectral methods for solving partial differential equations, *in* A. Iserles (ed.), *Acta Numerica*, Cambridge University Press, New York, pp. 203–267.

Fornberg, B. and Whitham, G. B.: 1978, A numerical and theoretical study of certain nonlinear wave phenomena, *Philosophical Transactions of the Royal Society of London* **289**, 373–404. Develops efficient Fourier pseudospectral method for Korteweg–deVries and related wave equations; the linear terms, which have constant coefficient, are integrated in time exactly.

Foster, I. T. and Worley, P. H.: 1997, Parallel algorithms for the spectral transform method, *SIAM J. Sci. Comput.* **18**(3), 806–837. Comparison of methods for spherical harmonics transforms for the nonlinear shallow water wave equations on various parallel architectures.

Fox, D. G. and Orszag, S. A.: 1973, Pseudospectral approximation to two-dimensional turbulence, *Journal of Computational Physics* **11**, 612–619.

Fox, L.: 1962, Chebyshev methods for ordinary differential equations, *Computer Journal* **4**, 318–331.

Fox, L. and Parker, I. B.: 1968, *Chebyshev Polynomials in Numerical Analysis*, Oxford University Press, London. Very readable and still good for general background, but the Clenshaw-type recursive algorithms for solving differential and integral equations are no longer popular except for special applications.

Fox, L., Hayes, L. and Mayers, D. F.: 1973, The double eigenvalue problem, *in* J. C. P. Miller (ed.), *Numerical Analysis*, Academic Press, pp. 93–112. Chebyshev solution of a 2d order ODE which has three boundary conditions imposed. Solutions exist because the problem has two independent eigenparameters.

Francis, P. E.: 1972, The possible use of Laguerre polynomials for representing the vertical structure of numerical models of the atmosphere, *Quart. J. Roy. Met. Soc.* **98**, 662–667. See also the correspondence by B. J. Hoskins and Francis, QJRMS 99, 571–572. The conclusion is that Laguerre polynomials demand a very short timestep, and are therefore useless in this context.

Francken, P., Deville, M. O. and Mund, E. H.: 1990, On the spectrum of the iteration operator associated to the finite element preconditioning of Chebyshev collocation calculations, *Computer Methods in Applied Mechanics and Engineering* **80**, 295–304.

Fraser, W. and Wilson, M. W.: 1966, Remarks on the Clenshaw-Curtis quadrature scheme, *SIAM Review* **8**, 322–327.

Froes Bunchaft, M. E.: 1997, Some extensions of the Lanczos-Ortiz theory of canonical polynomials in the Tau method, *Mathematics of Computation* **66**(218), 609–621. Extends and simplifies the canonical polynomial tau method; no numerical illustrations.

Frutos, J. and Sanz-Serna, J. M.: 1992, An easily implementable fourth-order method for the time integration of wave problems, *Journal of Computational Physics* **103**, 160–168.

Frutos, J. et al.: 1990, A Hamiltonian explicit algorithm with spectral accuracy for the 'good' Boussinesq equation, *Computer Methods in Applied Mechanics and Engineering* **80**, 417–423.

Funaro, D.: 1986, A multidomain spectral approximation of elliptic equations, *Methods for Partial Differential Equations* **2**, 187–205.

Funaro, D.: 1987a, Some results about the spectrum of the Chebyshev differencing operator, *in* E. L. Ortiz (ed.), *Numerical Approximations of P. D. E., Part III*, North-Holland, Amsterdam, pp. 271–284.

Funaro, D.: 1987b, A preconditioning matrix for the Chebyshev differencing operator, *SIAM Journal of Numerical Analysis* **24**, 1024–1031.

Funaro, D.: 1988a, Computing the inverse of the Chebyshev collocation derivative matrix, *SIAM Journal of Scientific and Statistical Computing* **9**, 1050–1058.

Funaro, D.: 1988b, Domain decomposition methods for pseudo spectral approximations. Part I. Second order equations in one dimension, *Numerische Mathematik* **52**, 329–344.

Funaro, D.: 1990a, Computational aspects of pseudospectral Laguerre approximations, *Appl. Numer. Math.* **6**(6), 447–457.

Funaro, D.: 1990b, Convergence analysis for pseudospectral multidomain approximations of linear advection equations, *IMA J. Numer. Anal.* **10**(1), 63–74.

Funaro, D.: 1990c, A variational formulation for the Chebyshev pseudospectral approximation of Neumann problems, *SIAM J. Numer. Anal.* **27**(3), 695–703.

Funaro, D.: 1991, Pseudospectral approximation of a PDE defined on a triangle, *Appl. Math. Comput.* **42**, 121–138. Subdivides each triangle into three quadrilaterals.

Funaro, D.: 1992a, Approximation by the Legendre collocation method of a model problem in electrophysiology, *Journal of Computational and Applied Mathematics* **43**, 261–271.

Funaro, D.: 1992b, *Polynomial Approximation of Differential Equations*, Springer-Verlag, New York. 313 pp.

Funaro, D.: 1993a, A new scheme for the approximation of advection-diffusion equations by collocation, *SIAM Journal of Numerical Analysis* **30**(6), 1664–1676.

Funaro, D.: 1993b, FORTRAN routines for spectral methods, *Report 891*, I. A. N.-C. N. R., Pavia, Italy. The 82 FORTRAN routines plus manual are available at the Web address ftp:/ftp.ian.pv.cnr.it/pub/splib.

Funaro, D.: 1994a, A fast solver for elliptic boundary-value problems in the square, *in* C. Bernardi and Y. Maday (eds), *Analysis, Algorithms and Applications of Spectral and High Order Methods for Partial Differential Equations*, Selected Papers from the International Conference on Spectral and High Order Methods (ICOSAHOM '92), Le Corum, Montpellier, France, 22-26 June 1992, North-Holland, Amsterdam, pp. 253–256. Also in Comp. Meths. Appl. Mech. Engrg., 116.

Funaro, D.: 1994b, Spectral elements in the approximation of boundary-value-problems in complex geometries, *Applied Numerical Mathematics* **15**(2), 201–205.

Funaro, D.: 1997a, Some remarks about the collocation method on a modified Legendre grid, *Computers and Mathematics with Applications* **33**, 95–103.

Funaro, D.: 1997b, *Spectral Elements for Transport-Dominated Equations*, Vol. 1 of *Lecture Notes in Computational Science and Engineering*, Springer-Verlag, Heidelberg. 200 pp.

Funaro, D. and Gottlieb, D.: 1988, A new method of imposing boundary conditions in pseudospectral approximations of hyperbolic equations, *Mathematics of Computation* **51**, 599–613.

Funaro, D. and Gottlieb, D.: 1991, Convergence results for pseudospectral approximations of hyperbolic systems by a penalty-type boundary treatment, *Mathematics of Computation* **57**, 585–596.

Funaro, D. and Heinrichs, W.: 1990, Some results about the pseudospectral approximation of one dimensional fourth order problems, *Numerische Mathematik* **58**, 399–418.

Funaro, D. and Kavian, O.: 1991, Approximation of some diffusion evolutions equations in unbounded domains by Hermite functions, *Mathematics of Computation* **57**, 597–619.

Funaro, D. and Rothman, E.: 1989, Preconditioning matrices for the pseudospectral approximations of first-order operators, *in* T. J. Chung and G. R. Karr (eds), *Finite Elements Analysis in Fluids*, UAH Press, Huntsville, Alabama, pp. 1458–1463.

Funaro, D. and Russo, A.: 1993, Approximation of advection-diffusion problems by a modified Legendre grid, *in* K. Morgan, E. Onate, J. Periaux, J. Peraire and O. C. Zienkiewicz (eds), *Finite Elements in Fluids, New Trends and Applications*, Pineridge Press, Battersea, England, pp. 1311–1318.

Funaro, D., Quarteroni, A. and Zanolli, P.: 1988, An iterative procedure with interface relaxation for domain decomposition methods, *SIAM Journal of Numerical Analysis* **25**, 1213–1236.

Funaro, O. C. D. and Kavian, O.: 1990, Laguerre spectral approximations of elliptic problems in exterior domains, *Computer Methods in Applied Mechanics and Engineering* **80**, 451–458.

Gad-el-Hak, M., Davis, S. H., McMurray, J. T. and Orszag, S. A.: 1984, On the stability of the decelerating laminar boundary layer, *Journal of Fluid Mechanics* **138**, 297–323.

Garba, A.: 1998, A mixed spectral/wavelet method for the solution of Stokes problem, *J. Comput. Phys.* **145**(1), 297–315. Chebyshev polynomials in the non-periodic coordinate, Daubechies wavelets in the spatially periodic coordinate.

García-Archilla, B.: 1995, Some practical experience with the time integration of dissipative equations, *Journal of Computational Physics* **122**, 25–29. Kuramoto-Sivashinsky equation is used to compare Nonlinear Galerkin and standard Galerkin methods; finds that standard stiff ODE methods are very effective, more so than Nonlinear Galerkin schemes.

García-Archilla, B.: 1996, A spectral method for the equal width equation, *Journal of Computational Physics* **125**, 395–402.

García-Archilla, B. and de Frutos, J.: 1995, Time integration of the non-linear Galerkin method, *IMA Journal of Numerical Analysis* **15**, 221–244. "The results show that for these problems [Kuramoto-Sivashinsky and reaction-diffusion equations], the nonlinear Galerkin method is not competitive with either pure spectral Galerkin or pseudospectral discretizations", from the abstract.

Gardner, D. R., Trogdon, S. A. and Douglass, R. W.: 1989, A modified tau spectral method that eliminates spurious eigenvalues, *Journal of Computational Physics* **80**, 137–167.

Gary, J. and Helgason, R.: 1970, A matrix method for ordinary differential equation eigenvalue problems, *Journal of Computational Physics* **5**, 169–187. Shows that the QR and QZ matrix eigensolvers make high order discretizations especially advantageous.

Gautheir, S., Guillard, H., Lumpp, T., Malé, J., Peyret, R. and Renaud, F.: 1996, A spectral domain decomposition technique with moving interfaces for viscous incompressible flows, *ECCOMAS 96*, John Wiley, New York.

Gelb, A.: 1997, The resolution of the Gibbs phenomenon for spherical harmonics, *Math. Comput.* **66**, 699–717. Applies the Gegenbauer polynomial sequence acceleration of Gottlieb and Shu to discontinuous functions on the sphere.

Gelb, A. and Gottlieb, D.: 1998, Recovering grid-point values without Gibbs oscillations in two dimensional domains on the sphere, *J. Comput. Phys.* Submitted. Gegenbauer polynomial regularization for rectangular regions or a union of rectancles on the sphere with meteorological applications.

Gentleman, W. M.: 1972a, Implementing Clenshaw-Curtis quadrature, *Communications of the ACM* **15**(5), 353–355. Table with complete FORTRAN code including the necesary cosine-FFT.

Gentleman, W. M.: 1972b, Implementing Clenshaw-Curtis quadrature: I. Methodology and experience, *Communications of the ACM* **15**(5), 337–342. Careful analysis of performance on fifty test integrals.

Gentleman, W. M.: 1972c, Implementing Clenshaw-Curtis quadrature: II. Computing the cosine transformation, *Communications of the ACM* **15**(5), 343–346.

Ghaddar, N. K., Korczak, K. Z., Mikic, B. B. and Patera, A. T.: 1986a, Numerical investigation of incompressible flow in grooved channels, Part 1: Stability and self-sustained oscillations, *Journal of Fluid Mechanics* **163**, 99–127.

Ghaddar, N. K., Korczak, K. Z., Mikic, B. B. and Patera, A. T.: 1986b, Numerical investigation of incompressible flow in grooved channels, Part 2: Resonance and oscillatory heat transfer, *Journal of Fluid Mechanics* **168**, 541–567.

Ghosh, S., Hossain, M. and Matthaeus, W. H.: 1993, The application of spectral methods in simulating compressible fluid and magnetofluid turbulence, *Computer Physics Communications* **74**, 18–40.

Gill, A. W. and Sneddon, G. E.: 1995, Complex mapped matrix methods in hydrodynamic stability problems, *Journal of Computational Physics* **122**, 13–24. Derive analytic formulas for optimizing Boyd's mappings to detour around singularities in the complex plane to calculate eigenvalues for modes with a singularity on or near the real axis, as occur in linearized hydrodynamic stability and Sturm-Liouville eigenproblems of the Fourth Kind; illustrated with experiments.

Gill, A. W. and Sneddon, G. E.: 1996, Pseudospectral methods and composite complex maps for near-boundary critical latitudes, *Journal of Computational Physics* **129**(1), 1–7. Simple formulas for optimizing a change-of-coordinate to resolve a differential equation with singularities very close to the boundary; Chebyshev pseudospectral solution of eigenproblems.

Giraldo, F. X.: 1998, The Lagrange-Galerkin spectral element method on unstructured quadrilateral grids, *J. Comput. Phys.* **147**(1), 114–146. Semi-Lagrangian spectral elements for fluids.

Givi, P. and Madnia, C. K.: 1993, Spectral methods in combustion, *in* T. J. Chung (ed.), *Numerical Modeling in Combustion*, Hemisphere, Taylor and Francis, Washington, pp. 409–452.

Glatzmaier, G. A.: 1984, Numerical simulations of stellar convective dynamos. Part I. The model and method, *Journal of Computational Physics* **55**, 461–484. Spherical harmonics in latitude and longitude, Chebyshev polynomials in radius. This model, with refinements, has been used in more than forty successive articles on mantle convection and stellar fluid dynamics by Glatzmaier and collaborators.

Glatzmaier, G. A.: 1988, Numerical simulations of mantle convection: Time-dependent ,three-dimensional, compressible spherical shell, *Geophysical and Astrophysical Fluid Dynamics* **43**, 223–264.

Glatzmaier, G. A. and Roberts, P. H.: 1997, Simulating the geodynamo, *Contemp. Phys.* **38**(4), 269–288. C.

Glowinski, R., Keller, H. B. and Reinhart, L.: 1985, Continuation conjugate gradient methods for the least squares solution of nonlinear boundary value problems, *SIAM Journal of Scientific and Statistical Computing* **6**, 793–832.

Godon, P.: 1995, The propagation of acoustic waves and quasi-periodic oscillations in accretion disc boundary layers, *Monthly Notices of the Royal Astronomical Society* **274**, 61–74. Polar coordinates in a two-dimensional plane; Fourier-Chebyshev with 4th order Runge-Kutta. Application.

Godon, P.: 1996a, Accretion disc boundary layers around pre-main-sequence stars, *Monthly Notices of the Royal Astronomical Society* **279**(4), 1071–1082. Polar coordinates in a two-dimensional plane; Fourier-Chebyshev with 4th order Runge-Kutta. Application.

Godon, P.: 1996b, Accretion disk boundary layers in classical t tauri stars, *Astrophysical Journal* **463**(2), 674–680. Polar coordinates in a two-dimensional plane; Fourier-Chebyshev with 4th order Runge-Kutta. Kosloff/Tal-Ezer mapping.

Godon, P.: 1997a, Advection in accretion disk boundary layers, *Astrophysical Journal* **483**(1), 882–886. Polar coordinates in a two-dimensional plane through the center of an accretion disk around a star: Chebyshev in radius and Fourier in the polar angle.

Godon, P.: 1997b, Numerical modeling of tidal effects in polytropic accretion disks, *Astrophysical Journal* **480**(1), 329–343. Polar coordinates in a two-dimensional plane through the center of an accretion disk around a star: Chebyshev in radius and Fourier in the polar angle. Kosloff-Tal-Ezer mapping.

Godon, P., , Regev, O. and Shaviv, G.: 1995, One-dimensional time-dependent numerical modeling of accretion disc boundary-layers, *Monthly Notices of the Royal Astronomical Society* **275**(4), 1093–1101. Chebyshev-Fourier polar coordinate model with 4th order Runge-Kutta time-marching.

Godon, P. and Shaviv, G.: 1993, A two-dimensional time-dependent Chebyshev method of collocation for the study of astrophysical flows, *Comput. Meths. Appl. Mech. Engin.* **110**(1–2), 171–194. Chebyshev-Fourier polar coordinate model with 4th order Runge-Kutta time-marching.

Godon, P. and Shaviv, G.: 1995, The dynamics of two-dimensional local and finite perturbations in envelopes of rotating dwarf stars, *Astrophysical Journal* **447**, 797–806. Chebyshev-Fourier polar coordinate model with 4th order Runge-Kutta time-marching.

Goldhirsch, I., Orszag, S. A. and Maulik, B. K.: 1987, An efficient method for computing leading eigenvalues and eigenvectors of large asymmetric matrices, *Journal of Scientific Computing* **2**, 33–58.

Gordon, C. T. and Stern, W. F.: 1982, A description of the GFDL global spectral model, *Monthly Weather Review* **110**, 625–644.

Gottlieb, D.: 1981, The stability of pseudospectral Chebyshev methods, *Mathematics of Computation* **36**, 107–118.

Gottlieb, D.: 1984, Spectral methods for compressible flow problems, *in* Soubbaramayer and J. P. Boujot (eds), *Proceedings of the 9th International Conference on Numerical Methods in Fluid Dynamics, Saclay, France*, number 218 in *Lecture Notes in Physics*, Springer-Verlag, New York, pp. 48–61.

Gottlieb, D. and Hirsh, R. S.: 1989, Parallel pseudospectral domain decomposition techniques, *Journal of Scientific Computing* **4**(1), 309–326.

Gottlieb, D. and Lustman, L.: 1983a, The Dufort-Frankel Chebyshev method for parabolic initial value problems, *Computers and Fluids* **11**, 107–120.

Gottlieb, D. and Lustman, L.: 1983b, The spectrum of the Chebyshev collocation operator for the heat equation, *SIAM Journal of Numerical Analysis* **20**, 909–921. Global and subdomain cases are analyzed with numerical results for the potential equation of airfoil theory.

Gottlieb, D. and Orszag, S. A.: 1977, *Numerical Analysis of Spectral Methods*, SIAM, Philadelphia, PA. 200 pp.

Gottlieb, D. and Orszag, S. A.: 1980, High resolution spectral calculations of inviscid compressible flows, *in* (ed.), *Approximation Methods for Navier-Stokes Problems*, Springer-Verlag, New York, pp. 381–398.

Gottlieb, D. and Shu, C.-W.: 1994, Resolution properties of the Fourier method for discontinuous waves, *in* C. Bernardi and Y. Maday (eds), *Analysis, Algorithms and Applications of Spectral and High Order Methods for Partial Differential Equations*, Selected Papers from the International Conference on Spectral and High Order Methods (ICOSAHOM '92), Le Corum, Montpellier, France, 22-26 June 1992, North-Holland, Amsterdam, pp. 27–38. Also in Comput. Meths. Appl. Mech. Engrg., vol. 116.

Gottlieb, D. and Shu, C.-W.: 1995a, On the Gibbs phenomenon IV: Recovering exponential accuracy in a subinterval from a Gegenbauer partial sum of a piecewise analytic function, *Mathematics of Computation* **64**, 1081–1095.

Gottlieb, D. and Shu, C.-W.: 1995b, On the Gibbs phenomenon V: Recovering exponential accuracy from collocation point values of a piecewise analytic function, *Numerische Mathematik* **71**, 511–526.

Gottlieb, D. and Streett, C. L.: 1990, Quadrature imposition of compatibility conditions in Chebyshev methods, *Journal of Scientific Computing* **5**(3), 223–240.

Gottlieb, D. and Tadmor, E.: 1985, Recovering pointwise values of discontinuous data within spectral accuracy, *in* E. M. Murman and S. S. Abarbanel (eds), *Progress and Supercomputing in Computational Fluid Dynamics*, Birkhäuser, Boston, pp. 357–375.

Gottlieb, D. and Tadmor, E.: 1991, The CFL condition for spectral approximation to hyperbolic BVPs, *Mathematics of Computation* **56**, 565–588.

Gottlieb, D. and Témam, R.: 1993, Implementation of the Nonlinear Galerkin method with pseudospectral (collocation) discretizations, *Applied Numerical Mathematics* **12**, 119–134.

Gottlieb, D. and Turkel, E.: 1980, On time discretization for spectral methods, *Studies in Applied Mathematics* **63**, 67–86.

Gottlieb, D. and Turkel, E.: 1985, Topics in spectral methods for time dependent problems, *in* F. Brezzi (ed.), *Numerical Methods In Fluid Dynamics*, Springer-Verlag, New York, pp. 115–155.

Gottlieb, D., Hussaini, M. Y. and Orszag, S. A.: 1984a, Theory and application of spectral methods, *in* R. G. Voigt, D. Gottlieb and M. Y. Hussaini (eds), *Spectral Methods for Partial Differential Equations*, SIAM, Philadelphia, pp. 1–54.

Gottlieb, D., Lustman, L. and Orszag, S. A.: 1981a, Spectral calculations of one-dimensional inviscid compressible flow, *SIAM Journal of Scientific and Statistical Computing* **2**, 296–310.

Gottlieb, D., Lustman, L. and Streett, C. L.: 1984b, Spectral methods for two-dimensional shocks, *in* R. G. Voigt, D. Gottlieb and M. Y. Hussaini (eds), *Spectral Methods for Partial Differential Equations*, SIAM, Philadelphia, pp. 79–96.

Gottlieb, D., Lustman, L. and Tadmor, E.: 1987a, Stability analysis of spectral methods for hyperbolic initial-value problems, *SIAM Journal of Numerical Analysis* **24**, 241–258.

Gottlieb, D., Lustman, L. and Tadmor, E.: 1987b, Convergence of spectral methods for hyperbolic initial-value problems, *SIAM Journal of Numerical Analysis* **24**, 532–537.

Gottlieb, D., Orszag, S. A. and Turkel, E.: 1981b, Stability of pseudospectral and finite difference methods for variable coefficient problems, *Mathematics of Computation* **37**, 293–305.

Gottlieb, D., Shu, C.-W., Solomonoff, A. and Vandeven, H.: 1992, On the Gibbs phenomenon I: recovering exponential accuracy from the Fourier partial sum of a nonperiodic analytic function, *Journal of Computational and Applied Mathematics* **43**, 81–98.

Grant, F. C. and Feix, M. R.: 1967, Fourier-Hermite solutions of the Vlasov equations in the linearized limit, *The Physics of Fluids* **10**(4), 696–702. Damping term is added to improve the otherwise slow convergence of the Hermite series for the velocity coordinate.

Gravel, S. and Staniforth, A.: 1994, A mass-conserving semi-Lagrangian scheme for shallow-water equations, *Monthly Weather Review* **122**, 243–248.

Gravel, S., Staniforth, A. and Côté, J.: 1993, A stability analysis of a family of baroclinic semi-Lagrangian forecast models, *Monthly Weather Review* **121**, 815–824.

Greengard, L. and Strain, J.: 1991, The fast Gauss transform, *SIAM J. Scient. Stat. Comput.* **12**, 79–94. Sum of N Gaussians at N points in space; not a generalized FFT but a close cousin.

Gresho, P. M., Gartling, D. K., Torczynski, J. R., Cliffe, K. A., Winters, K. H., Garratt, T. J., Spence, A. and Goodrich, J. W.: 1993, Is the steady visous incompressible two-dimensional flow over a backward- facing step at Re=800 stable?, *International Journal for Numerical Methods in Fluids* **17**, 501–541. Shows, by applying several different algorithms to a classical benchmark, that a published spectral element calculation was in error because of insufficient convergence tests. See also Kaiktsis, Karniadakis and Orszag(1991, 1996).

Grosch, C. E. and Orszag, S. A.: 1977, Numerical solution of problems in unbounded regions: coordinate transforms, *Journal of Computational Physics* **25**, 273–296.

Guillard, H. and Desideri, J. A.: 1990, Iterative methods with spectral preconditioning for elliptic equations, *Comput. Meths. Appl. Mech. Engrg.* **80**(1–3), 305–312.

Guillard, H. and Peyret, R.: 1988, On the use of spectral methods for the numerical solution of stiff problems, *Computer Methods in Applied Mechanics and Engineering* **66**, 17–43. Flame propagation problems and Burgers' equation.

Guo, B.-Y.: 1998, *Spectral Methods and Their Applications*, World Scientific, Singapore. 360 pp. Emphasizes proofs rather than numerical examples.

Guo, B.-Y. and Manoranjan, V. S.: 1985, A spectral method for solving the RLW equation, *IMA Journal of Numerical Analysis* **5**, 307–318.

Gustafsson, N. and McDonald, A.: 1996, A comparison of the HIRLAM gridpoint and spectral semi-Lagrangian models, *Monthly Weather Review* **124**(9), 2008–2022. Both models work well and at comparable cost; semi-Lagrangian advection is more accurate than the alternatives.

Gwynllyw, D. R., Davies, A. R. and Phillips, T. N.: 1996, A moving spectral element approach to the dynamically loaded journal bearing problem, *Journal of Computational Physics* **123**, 476–494.

Haidvogel, D. B.: 1977, Quasigeostrophic regional and general circulation modelling: an efficient pseudospectral aproximation technique, *in* R. P. Shaw (ed.), *Computing Methods in Geophysical Mechanics, Volume 25*, ASME, New York. REVIEW.

Haidvogel, D. B.: 1983, Periodic and regional models, *in* A. P. Robinson (ed.), *Eddies in Marine Science*, Springer-Verlag, New York, pp. 404–437. REVIEW.

Haidvogel, D. B. and Zang, T. A.: 1979, The accurate solution of Poisson's equation by expansion in Chebyshev polynomials, *Journal of Computational Physics* **30**, 167–180. Classic paper on separable BVP PDE's.

Haj, A., Phillips, C. and Delves, L. M.: 1980, The global element method for stationary advective problems, *International Journal of Numerical Methods in Engineering* **15**, 167–175.

Hald, O. H.: 1981, Convergence of Fourier methods for Navier-Stokes equations, *Journal of Computational Physics* **40**, 305–317. Proofs.

Haldenwang, P., Labrosse, G., Abboudi, S. and Deville, M.: 1984, Chebyshev 3-d spectral and 2-d pseudospectral solvers for the Helmholtz equation, *Journal of Computational Physics* **55**, 115–128. Comparison of three methods for constant coefficient PDE: diagonalization in two dimensions plus solution of tridiagonal systems in the third (Haidvogel and Zang, 1979), diagonalization in all three dimensions, and an iteration preconditioned by a fast direct finite difference solver for a separable system of cost $O(N^3 \log_2(N))$. The HZ algorithm was fastest, but loses about five digits due to roundoff for N as small as 64.

Haltiner, G. J. and Williams, R. T.: 1980, *Numerical Prediction and Dynamic Meteorology*, second edn, John Wiley. Chapter on spherical harmonics.

Han, H. C., Schultz, W. W., Boyd, J. P. and Schumack, M. R.: 1999, The flow in a elliptical journal bearing by the spectral element method for a rotating and translating shaft. To appear. The movement of the spinning central shaft was calculated simultaneously with the flow.

Hardiker, V.: 1997, A global numerical weather prediction model with variable resolution, *Monthly Weather Review* **125**(1), 59–73. Conformal mapping is used to make at T-83 spherical harmonics model as effective as a T-170 uniform resolution code for modelling hurricanes.

Harding, R. C.: 1968, Response of a one-dimensional Vlasov plasma to external electric fields, *Physics of Fluids* **11**(10), 2233–2240. Fourier for space coordinate, Hermite for velocity coordinate, finite difference for time. Little about numerics except that several hundred Hermite modes were needed because of problems resolved by later improvements (Holloway, 1996a,b).

Hargittai, I. and Hargittai, M.: 1994, *Symmetry: A Unifying Concept*, Shelter Publications, Bolinas, California. Good popularization of symmetry in art and nature.

Haugen, J. E. and Machenhauer, B.: 1993, A spectral limited-area model formulation with time-dependent boundary conditions applied to the shallow-water equations, *Monthly Weather Review* **121**, 2618–2630.

Haupt, S. E. and Boyd, J. P.: 1988, Modeling nonlinear resonance: A modification to Stokes' perturbation expansion, *Wave Motion* **10**, 83–98. Fourier basis for nonlinear eigenproblem. Analysis of the relationship between perturbation theory and the Galerkin numerical algorithm.

Haupt, S. E. and Boyd, J. P.: 1991, Double cnoidal waves of the Korteweg-deVries equation: The boundary value approach, *Physica D* **50**, 117–134. Two-dimensional Fourier basis for a nonlinear problem with two eigenparameters.

Haurwitz, B.: 1940, The motion of atmospheric disturbances on the spherical earth, *J. Marine Res.* **3**, 254–267. Shows spherical harmonics are travelling wave solutions of the shallow water equation, now known as "Rossby-Haurwitz" waves.

Healy, D. M., Rockmore, D. N., Kostelec, P. and Moore, S. S. B.: 1999, FFTs for the 2-sphere — Improvements and variations, *Adv. Appl. Math.* Submitted. Fast spherical harmonic transforms.

Heikes, R. and Randall, D. A.: 1995a, Numerical integration of the shallow-water equations on a twisted icosahedral grid. Part I: Basic design and result of tests., *Monthly Weather Review* **123**, 1862–1880. Non-spectral alternative to spherical harmonics.

Heikes, R. and Randall, D. A.: 1995b, Numerical integration of the shallow-water equations on a twisted icosahedral grid. Part II: A detailed description of the grid and an analysis of numerical accuracy., *Monthly Weather Review* **123**, 1881–1887. Non-spectral alternative to spherical harmonics.

Heinrichs, W.: 1987, *Kollokationsverfahren und Mehrgittermethoden bei elliptschen Randwertaufgaben*, Vol. 168 of *GMD-Bericht Nr.*, Oldenbourg-Verlag, Oldenbourg. Doctoral thesis.

Heinrichs, W.: 1988a, Line relaxation for spectral multigrid, *Journal of Computational Physics* **77**, 166–182.

Heinrichs, W.: 1988b, Multigrid methods for combined finite difference and Fourier problems, *Journal of Computational Physics* **78**, 424–436.

Heinrichs, W.: 1988c, Collocation and full multigrid methods, *Applied Mathematics and Computation* **28**, 35–45.

Heinrichs, W.: 1989a, Improved condition number for spectral methods, *Mathematics of Computation* **53**, 103–119.

Heinrichs, W.: 1989b, Spectral methods with sparse matrices, *Numerische Mathematik* **56**, 25–41.

Heinrichs, W.: 1989c, Konvergenzaussagen für Kollokationsverfahren bei elliptschen Randwertaufgaben, *Numerische Mathematik* **54**, 619–637.

Heinrichs, W.: 1990, Algebraic spectral multigrid methods, *Computer Methods in Applied Mechanics and Engineering* **80**, 281–289.

Heinrichs, W.: 1991a, A 3D spectral multigrid method, *Applied Mathematics and Computation* **41**, 117–128.

Heinrichs, W.: 1991b, Stabilization techniques for spectral methods, *Journal of Scientific Computing* **6**(1), 1–19. Basis functions that satisfy homogeneous boundary conditions reduce condition number.

Heinrichs, W.: 1991c, A stabilized treatment of the biharmonic operator with spectral methods, *SIAM Journal of Scientific and Statistical Computing* **12**, 1162–1172.

Heinrichs, W.: 1992a, A spectral multigrid method for the Stokes problem in stream-function formulation, *Journal of Computational Physics* **102**, 310–318.

Heinrichs, W.: 1992b, A stabilized multidomain approach for singular perturbation methods, *Journal of Scientific Computing* **7**, 95–127.

Heinrichs, W.: 1992c, Strong convergence estimates for pseudospectral methods, *Applications of Mathematics* **37**(6), 401–417.

Heinrichs, W.: 1992d, Spectral projective Newton-methods for quasilinear elliptic boundary value problems, *Calcolo* **29**, 33–48.

Heinrichs, W.: 1993a, Distributive relaxations for the spectral Stokes operator, *Journal of Scientific Computing* **8**, 389–398.

Heinrichs, W.: 1993b, Splitting techniques for the pseudospectral approximation of the unsteady Stokes equation, *SIAM Journal of Numerical Analysis* **30**, 19–39.

Heinrichs, W.: 1993c, Spectral multi-grid techniques for the Navier-Stokes equations, *Computer Methods in Applied Mechanics and Engineering* **106**, 297–314.

Heinrichs, W.: 1993d, Spectral multigrid methods for the reformulated Stokes equations, *Journal of Computational Physics* **107**(2), 213–224.

Heinrichs, W.: 1993e, Domain decomposition for fourth order problems, *SIAM Journal of Numerical Analysis* **30**(2), 435–453.

Heinrichs, W.: 1993f, Finite element preconditioning for spectral multigrid methods, *Applied Mathematics and Computation* **59**(1), 19–40.

Heinrichs, W.: 1993g, Spectral multigrid methods for domain decomposition problems using patching techniques, *Applied Mathematics and Computation* **59**(2), 165–176.

Heinrichs, W.: 1993h, Defect correction for convection dominated flow, *in* F.-K. Hebeker, R. Rannacher and G. Wittum (eds), *Proceedings of the International Workshop on Numerical Methods for the Navier-Stokes Equations*, number 47 in *Notes on Numerical Fluid Mechanics*, Heidelberg, pp. 111–121.

Heinrichs, W.: 1994a, Spectral methods for singular perturbation problems, *Applications of Mathematics* **39**(3), 161–188.

Heinrichs, W.: 1994b, *Efficient iterative solution of spectral systems for the Navier-Stokes equations*, Vol. 2 of *Wissenschaftliche Schriftenreihe Mathematik*, Verlag Dr. Koster, Berlin. Habilitationshrift, 109 pp.

Heinrichs, W.: 1994c, Spectral viscosity for convection dominated flow, *Journal of Scientific Computing* **9**(2), 137–147.

Heinrichs, W.: 1994d, Defection correction for the advection-diffusion equation, *Computer Methods in Applied Mechanics and Engineering* **119**, 191–197.

Heinrichs, W.: 1996, Defect correction for convection dominated flow, *SIAM Journal of Scientific Computing* **17**(5), 1082–1091.

Heinrichs, W.: 1998a, Spectral collocation on triangular elements, *J. Comput. Phys.* **145**(2), 743–757.

Heinrichs, W.: 1998b, Splitting techniques for the unsteady Stokes equations, *SIAM J. Numer. Anal.* **35**(4), 1646–1662. Third order time accurate Uzawa scheme with a single, unstaggered grid for in a pseudospectral approach.

Held, I. M. and Suarez, M. J.: 1994, A proposal for the intercomparison of the dynamical cores of atmospheric general circulation models, *Bull. Amer. Meteor. Soc.* **75**, 1825–1830. Comparison of 20-level models:, leapfrog time-stepping $T63$ spherical harmonics versus 2d order finite difference with $G72$[144 grid points around the equator]. There "is an impressive degree of agreement between the two models" but "the climate of both models are sensitive to resolution and have not yet converged at the resolution presented here". This article is unusual in claiming similar results for low order and high order methods for complex flows at the same resolution.

Hendry, J. A. and Delves, L. M.: 1979, The global element method applied to a harmonic mixed boundary value problem, *Journal of Computational Physics* **33**, 33–44.

Hendry, J. A., Delves, L. M. and Mohamed, J.: 1982, Iterative solution of the global element equations, *Computer Methods in Applied Mechanics and Engineering* **35**, 271–283.

Herbst, B. M. and Ablowitz, M.: 1992, Numerical homoclinic instabilities in the sine-Gordon equation, *Quaestiones Mathematicae* **15**, 345–363.

Herbst, B. M. and Ablowitz, M.: 1993, Numerical chaos, symplectic integrators, and exponentially small splitting distances, *Journal of Computational Physics* **105**, 122–132.

Herring, J. R., Orszag, S. A., Kraichnan, R. H. and Fox, D. G.: 1974, Decay of two-dimensional homogeneous turbulence, *Journal of Fluid Mechanics* **66**, 417–444.

Hesthaven, J. S.: 1998a, From electrostatics to almost optimal nodal sets for polynomial interpolation in a simplex, *SIAM Journal of Numerical Analysis* **35**(2), 655–676. Good distribution of points for a non-tensor product grid in a triangular subdomain. An alternative is given by Taylor&Wingate(1999).

Hesthaven, J. S.: 1998b, Integration preconditioning of pseudospectral operators. I. Basic linear operators, *SIAM J. Numer. Anal.* **35**(2), 1571–1593.

Hesthaven, J. S.: 1999, A stable penalty method for the compressible Navier-Stokes equations. III. Multidimensional domain decomposition schemes, *SIAM Journal of Scientific Computing* **20**(1), 62–93.

Hesthaven, J. S. and Gottlieb, D.: 1996, A stable penalty method for the compressible Navier-Stokes equations. I. Open boundary conditions., *SIAM Journal of Scientific Computing* **17**(3), 579–612.

Hesthaven, J. S. and Gottlieb, D.: 1999, Stable spectral methods for conservation laws on triangles with unstructured grids, *Comput. Meths. Appl. Mech. Engin.* In press.

Higgins, J. R.: 1977, *Completeness and Basis Properties of Sets of Special Functions*, Cambridge University Press, New York. This slim book is a very readable introduction to basis sets, but has no numerical content. Pp. 59-64 are a good discussion of complex-valued rational Chebyshev functions.

Hille, E.: 1939, Contributions to the theory of Hermitian series, *Duke Math. Journal* **5**, 875–936. This and the next three papers prove Hermite function convergence theorems.

Hille, E.: 1940a, Contributions to the theory of Hermitian series. II. The representation problem, *Transaction of the American Mathematical Society* **47**, 80–94.

Hille, E.: 1940b, A class of differential operators of infinite order, I, *Duke Mathematical Journal* **7**, 458–495.

Hille, E.: 1961, Sur les fonctions analytiques définies par des séries d'hermite, *Journal of Mathematiques Pures Appliques* **40**, 335–342.

Ho, L.-W. and Patera, A. T.: 1990, A Legendre spectral element method for simulation of unsteady incompressible viscous free-surface flows, *in* C. Canuto and A. Quarteroni (eds), *Spectral and High Order Methods for Partial Differential Equations: Proceedings of the ICOSAHOM '89 Conference in Como, Italy*, North-Holland/Elsevier, Amsterdam, pp. 355–366. Also in Comput. Meths. Appl. Mech. Engrg., vol. 80, with the same page numbers.

Ho, L. W. and Patera, A. T.: 1991, Variational formulation of three-dimensional viscous free-surface flows: natural imposition of surface tension boundary conditions, *International Journal for Numerical Methods in Fluids* **13**, 691–698.

Ho, L.-W. and Rønquist, E. M.: 1994, Spectral element solution of steady incompressible viscous free-surface flows, *in* C. Bernardi and Y. Maday (eds), *Analysis, Algorithms and Applications of Spectral and High Order Methods for Partial Differential Equations*, Selected Papers from the International Conference on Spectral and High Order Methods (ICOSAHOM '92), Le Corum, Montpellier, France, 22-26 June 1992, North-Holland, Amsterdam, pp. 347–368. Also in Finite Elements in Analysis and Design, vol. 16, pp. 207-229.

Ho, L.-W., Maday, Y., Patera, A. T. and Rønquist, E. M.: 1990, A high-order Lagrangian-decoupling method for the incompressible Navier-Stokes equations, *in* C. Canuto and A. Quarteroni (eds), *Spectral and High Order Methods for Partial Differential Equations: Proceedings of the ICOSAHOM '89 Conference in Como, Italy*, North-Holland/Elsevier, Amsterdam, pp. 65–90.

Hogan, T. F. and Rosmond, T. E.: 1991, The description of the Navy operational global atmospheric prediction systems's spectral forecast model, *Monthly Weather Review* **119**, 1786–1815.

Holloway, J. P.: 1996a, Spectral velocity discretizations for the Vlasov-Maxwell equations, *Transport Theory and Statistical Physics* **25**, 1–32.

Holloway, J. P.: 1996b, Hamiltonian spectral methods, *Journal of Computational Physics* **129**(1), 121–133. Fourier-Hermite spectral methods.

Holly, Jr., F. M. and Preissman, A.: 1977, Accurate evaluation of transport in two dimensions, *Journal of Hydraulics Division of the ASCE* **98**, 1259–1277. Semi-Lagrangian advection.

Holvorcem, P. R.: 1992, Asymptotic summation of Hermite series, *J. Phys. A* **25**(4), 909–924.

Holvorcem, P. R. and Vianna, M. L.: 1992, Integral-equation approach to tropical ocean dynamics. 2. Rossby-wave scattering from the equatorial Atlantic western boundary, *J. Marine Res.* **50**(1), 33–61. Boundary-element algorithm constructed through ingenious summation of slowly converging Hermite series.

Hortal, M. and Simmons, A. J.: 1991, Use of reduced Gaussian grids in spectral models, *Monthly Weather Review* **119**, 1057–1074.

Hoskins, B. J.: 1973, Comments on the possible use of Laguerre polynomials for representing the vertical structure of numerical models of the atmosphere, *Quart. J. Roy. Met. Soc.* **99**, 571–572.

Hoskins, B. J.: 1980, Representation of the earth topography using spherical harmonics, *Monthly Weather Review* **108**, 111–115. Exponential filter.

Hoskins, B. J. and Simmons, A. J.: 1975, A multi-layer spectral model and the semi-implicit method, *Quart. J. Roy. Met. Soc.* **101**, 637–655. Comparison of spherical harmonics/vertical finite differences model with a three-dimensional finite difference code.

Hua, B. L.: 1987, Periodic quasi-geostrophic models, *in* J. J. O'Brien (ed.), *Advanced Physical Oceanographical Numerical Modelling*, D. Reidel, Dordrect, Holland, pp. 233–254. REVIEW.

Hua, B. L. and Haidvogel, D. B.: 1986, Numerical simulations of the vertical structure of quasi-geostrophic turbulence, *Journal of the Atmospheric Sciences* **43**(23), 2923–2936. Fourier pseudospectral in the horizontal, normal modes in ocean depth for both constant and variable stratification.; detailed description of the numerical algorithms.

Huang, W.-Z. and Sloan, D. M.: 1993a, A new pseudospectral method with upwind features, *IMA Journal of Numerical Analysis* **13**, 413–430. The authors device is to approximate $u(x)$ by different polynomials in the same equation. Diffusive terms are treated by the usual Chebyshev-Lobatto approximation. The advective terms are approximated by a polynomial which omits one boundary point, keeping only the boundary value on the upwind side. When the shock layer is adequately resolved, upwindlng

is irrelevant. When the shock is underresolved. i. e., a boundary layer of thickness of ϵ where $\epsilon \ll l/N$, they show their scheme is much better behaved than the standard pseudospectral method. There is an $O(1)$ error around the shock, but the pollution of areas away from the shock is greatly reduced. They show many analogies between their scheme and upwinded finite differences.

Huang, W. Z. and Sloan, D. M.: 1993b, Pole condition for singular problems, *Journal of Computational Physics* **107**, 254. Spectral methods for polar and spherical coordinates.

Huang, W.-Z. and Sloan, D. M.: 1994, The pseudospectral method for solving differential eigenvalue equations, *Journal of Computational Physics* **111**, 399–409.

Hussaini, M. Y. and Zang, T. A.: 1984, Iterative spectral methods and spectral solution to compressible flows, *in* D. G. R. Voigt and M. Y. Hussaini (eds), *Spectral Methods for PDEs*, SIAM, Philadelphia.

Hussaini, M. Y., Kopriva, D. A. and Patera, A. T.: 1989, Spectral collocation methods, *Applied Numerical Mathematics* **5**, 177–208.

Hussaini, M. Y., Kopriva, D. A., Salas, M. D. and Zang, T. A.: 1985a, Spectral methods for the Euler equations: Part I — Fourier methods and shock-capturing, *AIAA J.* **23**, 64–70. Shows that spectral shock-capturing, that is, employing a smoothing technique to resolve a shock whose location is not directly calculated, fails for an astrophysical problem.

Hussaini, M. Y., Kopriva, D. A., Salas, M. D. and Zang, T. A.: 1985b, Spectral methods for the Euler equations: Part II – Chebyshev methods and shock-fitting, *AIAA J.* **23**, 234–240. Shock-fitting explicitly computes the location of the shock and then maps the computational domain so that the shock lies along a grid line in the new coordinate. This is much more effective than shock capturing for strong shocks.

If, F., Berg, P., Christiansen, P. L. and Skovgaard, O.: 1987, Split-step spectral method for nonlinear Schrodinger equation with absorbing boundaries, *Journal of Computational Physics* **72**, 501–503. Solve NLS equation with periodic boundary conditions on large period L to imitate the infinite interval. An artificial damping $\gamma(x)$ is added so that waves propagating left and right out of the computational zone do not return. The NLS equation is modified to $i\,u_t + \frac{1}{2}u_{xx} + |u|^2 u = -i\,\gamma(x)\,u$, $\gamma(x) = \gamma_0\{\text{sech}^2(\alpha[x - L/2]) + \text{sech}^2(\alpha[x + L/2])\}$ that is, the artificial damping is large at the ends of the interval, but exponentially small in the middle where the solution should resemble that of the undamped equation.

Ioakimidis, N. I.: 1987, Quadrature methods for the determination of zeros of transcendental functions — a review, *in* P. Keast and G. Fairweather (eds), *Numerical Integration: Recent Developments, Software and Applications*, D. Reidel, Dordrecht, Holland, pp. 61–82.

Ioakimidis, N. I. and Anastasselou, E. G.: 1986, An elementary noniterative quadrature-type method for the numerical solution of a nonlinear equation, *Computing* **37**, 269–275. Gives an explicit, exponentially convergent formula for the root of a function $f(x)$ within an interval by combining Chebyshev-Lobatto and Chebyshev-Gauss quadrature.

Iranzo, V. and Falques, A.: 1992, Some spectral approximations for differential equations in unbounded domains, *Computer Methods in Applied Mechanics and Engineering* **98**, 105–126. Compares Laguerre-tau, Laguerre pseudospectral, and rational Chebyshev pseudospectral for $x \in [0, \infty]$.

Iskandarani, M., Haidvogel, D. B. and Boyd, J. P.: 1994, A staggered spectral finite element method for the shallow water equations, *Int. J. Num. Meths. Fluids* **20**, 393–414.

Ito, K. and Teglas, R.: 1986, Legendre-Tau approximations for functional differential equations, *SIAM Journal for Control and Optimization* **24**, 737–759.

Ito, K. and Teglas, R.: 1987, Legendre-Tau approximations for functional differential equations, part 2: The linear quadratic optimal control problem, *SIAM Journal for Control and Optimization*.

Jackson, E., She, Z.-S. and Orszag, S. A.: 1991, A case study in parallel computing. I. Homogeneous turbulence on a hypercube, *Journal of Scientific Computing* **6**(1), 27–46.

Jacobs, S. J.: 1990, A variable order pseudospectral method for two-point boundary value problems, *Journal of Computational Physics* **88**, 169–182.

Jacobs, S. J.: 1995, An accurate split step scheme for viscous incompressible fluid flow, *Journal of Computational Physics* **119**, 26–33.

Jakob-Chien, R. and Alpert, B. K.: 1997, A fast spherical filter with uniform resolution, *Journal of Computational Physics* **136**(2), 580–584.

Jakob-Chien, R., Hack, J. J. and Williamson, D. L.: 1995, Spectral transform solutions to the shallow water test set, *Journal of Computational Physics* **119**, 164–187.

Jakob, R.: 1993, *Fast and Parallel Spectral Transform Algorithms for Global Shallow Water Models*, PhD dissertation, University of Colorado, Department of Electrical and Computer Engineering. Also available as NCAR Cooperative Thesis CT-144 from the National Center for Atmospheric Research, Publications Office, P. O. Box 3000,Boulder, CO 80307; author later known as R. Jakob-Chien.

Jarratt, M., Lund, J. and Bowers, K. L.: 1990, Galerkin schemes and the sinc-Galerkin method for singular Sturm-Liouville problems, *Journal of Computational Physics* **89**, 41–62.

Jarraud, M. and Baede, A. P. M.: 1985, The use of spectral techniques in numerical weather prediction, *in* B. Enquist, S. Osher and R. Somerville (eds), *Large Scale Computations in Fluid Mechanics*, number 22 in *Lectures in Applied Mathematics*, American Mathematical Society, Washington, pp. 1–41. REVIEW.

Jauberteau, F., Rosier, C. and Témam, R.: 1990a, A nonlinear Galerkin method for the Navier-Stokes equations, *Computer Methods in Applied Mechanics and Engineering* **80**, 245–260.

Jauberteau, F., Rosier, C. and Témam, R.: 1990b, The nonlinear Galerkin method in computational fluid dynamics, *Applied Numerical Mathematics* **6**, 361–370.

Jensen, T. G. and Kopriva, D. A.: 1989, Comparison of finite difference and spectral collocation reduced gravity ocean model, *in* A. M. Davies (ed.), *Modeling Marine Systems*, number II, CRC Press, Boca Raton, Florida, pp. 25–39.

Jolly, M. S. and Xiong, C.: 1995, On computing the long-time solution of the two-dimensional Navier-Stokes equations, *Theoretical and Computational Fluid Dynamics* **7**, 261–278. Comparison of several variants of the Nonlinear Galerkin method with standard two-dimensional Fourier method.

Jolly, M. S., Kevrekidis, I. G. and Titi, E. S.: 1990, Approximate inertial manifolds for the Kuramoto-Sivashinsky equation: Analysis and computations, *Physica D* **44**, 38–60. Low order Fourier basis.

Jolly, M. S., Kevrekidis, I. G. and Titi, E. S.: 1991, Preserving dissipation in approximate inertial forms for the Kuramoto-Sivashinsky equation, *Dynamics and Differential Equations* **3**(2), 179–197. Several variations of Nonlinear Galerkin methods.

Jones, D. A., Margolin, L. G. and Titi, E. S.: 1995, On the effectiveness of the approximate inertial manifold — a computational study, *Theoretical and Computational Fluid Dynamics* **7**, 243–260. Comparison of Nonlinear Galerkin with standard spectral Galerkin; mostly theory with numerical illustrations using forced Burgers equation and forced Kuramoto-Sivashinsky equation.

Jones, W. B. and O'Brien, J. J.: 1996, Pseudo-spectral methods and linear instabilities in reaction-diffusion fronts, *Chaos* **6**(2), 219–228. Comparisons of two-dimensional Fourier scheme with second order finite differences; Fourier is superior for both accuracy and preserving symmetries.

Joyce, G., Knorr, G. and Meier, H. K.: 1971, Numerical integration methods of the Vlasov equation, *Journal of Computational Physics* **8**, 53–63. Reviews Fourier ("characteristic function") and Hermite series and introduces a method of moments for the velocity coordinate. The poorly-converging coefficients of the moments expansion are truncated by polynomial extrapolation in degree n. (The method of moments, which is badly conditioned at high order, is *not* recommended.

Kaiktsis, L., Karniadakis, G. E. and Orszag, S. A.: 1991, Onset of three-dimensionality, equilibria, and early transition in flow over a backward facing step, *Journal of Fluid Mechanics* **231**, 501–528.

Kaiktsis, L., Karniadakis, G. E. and Orszag, S. A.: 1996, Unsteadiness and convective instabilities in two-dimensional flow over a backward facing step, *Journal of Fluid Mechanics* **321**, 157–187. Spectral elements. Attempt to understand why the authors' early paper gave unsteady flow whereas all other studies found only steady solutions.

Kalnay de Rivas, E.: 1972, On the use of nonuniform grids in finite-difference equations, *Journal of Computational Physics* **10**, 202–210. Coordinate mappings to resolve boundary layers. Not spectral, but her cosine mapping gives the usual Chebyshev pseudospectral grid.

Kanamitsu, M., Tada, K., Kudo, T., Sato, N. and Isa, S.: 1983, Description of the JMA operational spectral model, *Journal of the Meteorological Society of Japan* **61**, 812–828.

Karageorghis, A.: 1988a, A note on the Chebyshev coefficients of the general order derivative of an infinitely differentiable function, *Journal of Computational and Applied Mathematics* **21**, 129–132.

Karageorghis, A.: 1988b, A note on the Chebyshev coefficients of the moments of the general order derivative of an infinitely differentiable function, *Journal of Computational and Applied Mathematics* **21**, 383–386.

Karageorghis, A.: 1993, On the equivalence between basis recombination and boundary bordering formulations for spectral collocation methods in rectangular domains, *Mathematics and Computers in Simulation* **35**, 113–123.

Karageorghis, A.: 1994a, Satisfaction of boundary conditions for Chebyshev collocation methods in cuboidal domains, *Computers and Mathematics with Applications* **27**, 85–90.

Karageorghis, A.: 1994b, A conforming spectral technique for biharmonic-type problems in rectangular domains, *Journal of Computational and Applied Mathematics* **51**, 275–278.

Karageorghis, A.: 1994c, Conforming spectral methods for Poisson problems in cuboidal domains, *Mathematics and Computers in Simulation* **9**, 341–350.

Karageorghis, A.: 1995, A fully conforming spectral collocation scheme for second and fourth order problems, *Computer Methods in Applied Mechanics and Engineering* **126**, 305–314.

Karageorghis, A. and Phillips, T. N.: 1991, Conforming Chebyshev spectral collocation methods for the solution of laminar flow in a constricted channel, *IMA Journal of Numerical Analysis* **11**, 33–55.

Karageorghis, A. and Phillips, T. N.: 1992, On the coefficients of differentiated expansions of ultraspherical polynomials, *Applied Numerical Mathematics* **9**, 133–141.

Karagheorghis, A. and Paprzycki, M.: 1996a, Direct methods for spectral approximations in nonconforming domain decompositions, *Comput. Math. Appl.* **35**(11), 75–82.

Karagheorghis, A. and Paprzycki, M.: 1996b, An efficient direct method for fully conforming spectral collocation schemes, *Numerical Mathematics* **12**, 309–319. Solution of linear systems arising from fourth and second order problems through multidomain spectral methods.

Karniadakis, G. E. and Orszag, S. A.: 1993a, Some novel aspects of spectral methods, *in* M. Y. Hussaini, A. Kumar and M. D. Salas (eds), *Algorithmic Trends in Computational Fluid Dynamics*, Springer-Verlag, New York, p. 245.

Karniadakis, G. E. and Orszag, S. A.: 1993b, Nodes, modes and flow codes, *Physics Today* **46**, 32–42. REVIEW.

Karniadakis, G. E. and Sherwin, S. J.: 1995a, A triangular and tetrahedral basis for high-order finite elements: applications to the incompressible Navier-Stokes equations, *International Journal for Numerical Methods in Engineering* **38**, 3775–.

Karniadakis, G. E. and Sherwin, S. J.: 1995b, Atriangular spectral element method: applications to the incompressible Navier-Stokes equations, *Computer Methods in Applied Mechanics and Engineering* **123**, 189–229.

Karniadakis, G. E. and Sherwin, S. J.: 1999, *Spectral/hp Element Methods for CFD*, Oxford University Press, Oxford. 448 pp., 140 illustrations. Detailed analysis of both quadrilateral and triangular elements with many numerical examples.

Karniadakis, G. E., Bullister, E. T. and Patera, A. T.: 1986, A spectral element method for solution of two-and three-dimensional incompresssible Navier-Stokes equations, *in* P. Bergan and K. J. Bathe (eds), *Proceedings of the Europe-U.S. Conference on Finite Element Methods for Nonlinear Problems*, Wunderlich/Springer-Verlag, New York, pp. 803–817.

Karniadakis, G. E., Israeli, M. and Orszag, S. A.: 1991, High-order time-accurate splitting methods for incompressible Navier-Stokes equations, *Journal of Computational Physics* **97**, 414–443.

Kasahara, A.: 1977, Numerical integration of the global barotropic primitive equations with Hough harmonic expansions, *Journal of the Atmospheric Sciences* **34**, 687–701.

Kasahara, A.: 1978, Further studies on a spectral model of the global barotropic primitive equations with hough harmonic expansions, *Journal of the Atmospheric Sciences* **35**, 2043–2051.

Kasahara, A. and Puri, K.: 1981, Spectral representation of three-dimensional global data by expansion in normal modes, *Mon. Wea. Rev.* **109**, 37–51. Hough functions in latitude and longitude, eigenfunctions of the "vertical structure" equation in height.

Katopodes, N., Sanders, B. F. and Boyd, J. P.: 1998, Short wave behavior of long wave equations, *Waterways, Coastal and Ocean Engineering Journal of ASCE* **124**(5), 238–247. Application of Fourier basis; methods described in Sanders, Katopodes and Boyd (1998).

Keller, H. B.: 1977, Numerical solution of bifurcation and nonlinear eigenvalue problems, *in* P. Rabinowitz (ed.), *Applications of Bifurcation Theory*, Academic Press, New York, pp. 359–384.

Keller, H. B.: 1992, *Numerical Methods for Two-Point Boundary-Value Problems*, Dover, New York. Reprints of good research papers on solving nonlinear algebraic equations as well as differential equations.

Kermode, M., McKerrell, A. and Delves, L. M.: 1985, The calculation of singular coefficients, *Computer Methods in Applied Mechanics and Engineering* **50**, 205–215. Corner singularities, global elements.

Khabibrakhmanov, I. K. and Summers, D.: 1998, The use of generalized Laguerre polynomials in spectral methods for nonlinear differential equations, *Comput. Math. Appl.* **36**(2), 65–70. Galerkin method, implemented by recurrence, and illustrated by the Blasius equation on the semi-infinite domain.

Khajah, H. G. and Ortiz, E. L.: 1992, Numerical approximation of solutions of functional equations using the Tau method, *Applied Numerical Mathematics* **9**, 461–474.

Khajah, H. G. and Ortiz, E. L.: 1993, Rational approximations: a tau method approach, *in* T. M. Rassias, H. M. Srivastava and A. Yanushauskas (eds), *Topics in polynomials of one and several variables and their applications*, Volume dedicated to the memory of P. L. Chebyshev, World Scientific, Singapore, pp. 323–333.

Khorrami, M. R. and Malik, M. R.: 1993, Efficient computation of spatial eigenvalues for hydrodynamic stability analysis, *Journal of Computational Physics* **104**(1), 267–272. Show that by neglecting terms quadratic in the eigenvalue, one can halve the size of the matrix given the QZ algorithm, and then safely refine the eigenvalues by iterative correction with the quadratic term reinserted.

Kida, S.: 1985, Three-dimensional periodic flows with high-symmetry, *Journal of the Physical Society of Japan* **54**, 2132–2136. Three-dimensional Fourier basis, greatly reduced by symmetry.

Kida, S. and Orszag, S. A.: 1990, Energy and spectral dynamics in forced compressible turbulence, *Journal of Scientific Computing* **5**(2), 85–126. Fourier 64^3 with Runge-Kutta-Gill timestepping.

Kikuchi, N.: 1986, *Finite Element Methods in Mechanics*, Cambridge University Press, Cambridge. Very readable, elementary introduction to finite elements and skyline matrix solvers.

Klimas, A. J.: 1987, A method for overcoming the velocity space filamentation problem in collisionless plasma model solutions, *Journal of Computational Physics* **68**, 202–226. Fourier-Galerkin for both velocity and space coordinates; introduces a filter which is quite effective in controlling the dissipation of small-scale structure in the velocity coordinate.

Klimas, A. J. and Farrell, W. M.: 1994, A splitting algorithm for Vlasov simulation with filamentation filtration, *Journal of Computational Physics* **110**, 150–163. Fourier-Galerkin for both velocity coordinate and space coordinate; shows a particular filter greatly improves convergence and reduces filamentation in the velocity.

Kopriva, D. A.: 1986, A spectral multidomain method for the solution of hyperbolic systems, *Applied Numerical Mathematics* **2**, 221–241.

Kopriva, D. A.: 1987, A practical assessment of spectral accuracy for hyperbolic problems with discontinuity, *Journal of Scientific Computing* **2**, 249–262.

Kopriva, D. A.: 1988, A multidomain spectral collocation computation of the sound generated by a shock-vortex interaction, *in* M. Schultz, D. Lee and R. Sternberg (eds), *Computational Acoustics and Wave Propagation*, North-Holland, Amsterdam.

Kopriva, D. A.: 1989a, Computation of hyperbolic equations on complicated domains with patched and overset Chebyshev grids, *SIAM Journal of Scientific and Statistical Computing* **10**, 120–132.

Kopriva, D. A.: 1989b, Domain decomposition with both spectral and finite difference methods for the accurate computation of flows with shocks, *Applied Numerical Mathematics* **6**, 141–151.

Kopriva, D. A.: 1991, Multidomain spectral solution of the Euler gas dynamics equations, *Journal of Computational Physics* **96**, 428–450.

Kopriva, D. A.: 1992, Spectral solution of inviscid supersonic flows over wedges and axisymmetric cones, *Computers and Fluids* **21**, 247–266.

Kopriva, D. A.: 1993a, Spectral solution of the viscous blunt-body problem, *AIAA Journal* **31**, 1235–1242.

Kopriva, D. A.: 1993b, Spectral solutions of high-speed flows over blunt cones, *AIAA Journal* **31**, 2227–2231.

Kopriva, D. A.: 1993c, Multidomain spectral solution of compressible viscous flows, *AIAA Journal* **31**, 3376–3384.

Kopriva, D. A.: 1994, Multidomain spectral solution of compressible viscous flows, *Journal of Computational Physics* **115**(1), 184–199.

Kopriva, D. A.: 1996a, Spectral solution of the viscous blunt body problem. II. Multidomain approximation, *AIAA Journal* **34**(3), 560–564.

Kopriva, D. A.: 1996b, A conservative, staggered-grid multidomain method for the Euler gas-dynamics equations, *in* A. V. Ilin and L. R. Scott (eds), *Proceedings of the Third International Conference on Spectral and High Order Methods*, Houston Journal of Mathematics, Houston, Texas, pp. 457–468.

Kopriva, D. A.: 1998, A staggered-grid multidomain spectral method for the compressible Navier-Stokes equations, *Journal of Computational Physics* **143**(1), 125–158. Nonconforming interfaces; complex geometry.

Kopriva, D. A. and Hussaini, M. Y.: 1989, Multidomain spectral solution of shock-turbulence interactions, *in* T. F. Chan, R. Glowinski, J. Periaux and O. B. Widlund (eds), *Domain Decomposition Methods*, SIAM, Philadelphia, pp. 340–350.

Kopriva, D. A. and Kolias, J. H.: 1996, A conservative staggered-grid Chebyshev multidomain method for compressible flows, *Journal of Computational Physics* **125**, 244–261.

Kopriva, D. A. and Panchang, V. G.: 1989, Pseudospectral solution of two-dimensional water-wave propagation, *Mathematical and Computer Modelling* **12**, 625–640.

Kopriva, D. A., Zang, T. A. and Hussaini, M. Y.: 1991, Spectral methods for the Euler equations: The blunt body revisited, *AIAA Journal* **29**(9), 1458–1462.

Kopriva, D. A., Zang, T. A., Salas, M. D. and Hussaini, M. Y.: 1984, Pseudospectral solution of two-dimensional gas-dynamics problems, *in* M. Pandolfi and R. Piva (eds), *Proceedings 5th GAMM Conf. Numerical Methods on Fluid Mechanics*, Vieweg, Braunschweig, pp. 185–192.

Korczak, K. Z. and Patera, A. T.: 1986, An isoparametric spectral element method for solution of the Navier-Stokes equations in complex geometry, *Journal of Computational Physics* **62**, 361–382.

Kosloff, D., Reshef, M. and Loewenthal, D.: 1984, Elastic wave calculations by the Fourier method, *Bulletin of the Seismological Society of America* **74**, 875–891.

Kosloff, R. and Tal-Ezer, H.: 1993, A modified Chebyshev pseudospectral method with an $O(1/N)$ time step restriction, *Journal of Computational Physics* **104**, 457–469. The key idea is to employ a mapping which almost converts the Chebyshev polynomials back into cosines. The interpolahon grid is not uniform, but the roots of the mapped polynomials are not as concentrated near the endpoints as those of the Chebyshev polynomials. When the stretching parameter is chosen so as to give an $O(1/N)$ timstep, the asymptotic rate of convergence is reduced from geometric to algebraic, however.

Koures, V. G.: 1996, Solving the Coulomb Shrödinger equation in $d = 2 + 1$ via sinc collocation, *Journal of Computational Physics* **128**(1), 1–5.

Kreiss, H.-O.: 1991, Problems with different time scales, *Acta Numerica* **1**, 101–139. Numerical algorithms for problems with a "slow manifold".

Krishnamurti, T. N.: 1962, Numerical integration of primitive equations by a quasi-Lagrangian advection scheme, *Journal of Applied Meteorology* **1**, 508–521. Not spectral; early paper on semi-Lagrangian methods.

Kuo, H.-C. and Williams, R. T.: 1990, Semi-Lagrangian solutions to the inviscid Burgers equation, *Monthly Weather Review* **118**, 1278–1288. Elementary, but good introduction to semi-Lagrangian schemes.

Lacour, C. and Maday, Y.: 1997, Two different approaches for matching nonconforming grids: The mortar element method and the FETI method, *BIT* **37**(3), 720–738.

Lambiotte, J., Bokhari, S., Hussaini, M. Y. and Orszag, S. A.: 1982, Navier-Stokes solutions on the Cyber-203 by a pseudospectral technique, *10th IMACS World Congress on System Simulation and Scientific Computation*, Montreal. Splitting with mixed finite difference/spectral spatial discretization.

Lanczos, C.: 1938, Trigonometric interpolation of empirical and analytical functions, *Journal of Mathematics and Physics* **17**, 123–199. The origin of both the pseudospectral method and the tau method. Lanczos is to spectral methods what Newton was to calculus.

Lanczos, C.: 1956, *Applied Analysis*, Prentice-Hall, Englewood Cliffs, New Jersey. 400 pp.

Lanczos, C.: 1966, *Discourse on Fourier Series*, Oliver and Boyd, Edinburgh.

Lanczos, C.: 1973, Legendre versus Chebyshev polynomials, *in* J. C. P. Miller (ed.), *Numerical Analysis*, Academic Press, New York, pp. 191–201. Shows Legendre is "superconvergent" and much better than Chebyshev at $x = \pm 1$.

Lander, J. and Hoskins, B. J.: 1997, Believable scales and parameterizations in a spectral transform model, *Mon. Weath. Rev.* **125**, 292–303.

Laprise, R.: 1992, The resolution of global spectral methods, *Bulletin of the American Meteorological Society* **73**, 1453–1454.

Launay, J., Bouchet, S., Randriamampianina, A., Bontoux, P. and Gibart, P.: 1994, Modeling and experiments on epitaxial growth on a GaAs hemisphere substrate at 1 g and under hypergravity, *in* L. L. Regel and W. R. Wilcox (eds), *Materials Processing in High Gravity*, Plenum Press, New York, pp. 139–160. Two-dimensional (r, z) crystal in cylindrical coordinates using a Chebyshev tau method with the Haidvogel-Zang diagonalization method and time integration by the software package LSODA.

Le Quéré, P. and Pécheux, J.: 1989, Multiple transitions in axisymmetric annulus convection, *Journal of Fluid Mechanics* **206**, 517–544. Chebyshev-tau spectral method.

Le Quéré, P. and Pécheux, J.: 1990, A three-dimensional pseudo-spectral algorithm for the computation of convection in a rotating annulus, *Computer Methods in Applied Mechanics and Engineering* **80**, 261–271.

Lee, N. Y., Schultz, W. W. and Boyd, J. P.: 1989, Stability of fluid in a rectangular enclosure by spectral method, *International Journal of Heat and Mass Transfer* **32**, 513–520.

Leith, C. E.: 1980, Nonlinear normal mode initialization and quasi-geostrophic theory, *Journal of the Atmospheric Sciences* **37**, 958–968.

Leovy, C. B.: 1964, Simple models of thermally driven mesospheric circulations, *Journal of the Atmospheric Sciences* **21**, 327–341. Spectral eigensolution of Laplace's Tidal equation.

Leslie, L. M. and Purser, R. J.: 1991, High order numerics in an unstaggered three-dimensional time-split semi-Lagrangian forecast model, *Monthly Weather Review* **119**, 1612–1623. Not spectral, but very high order spatial differences.

Levin, J. G., Iskandarani, M. and Haidvogel, D. B.: 1997, A spectral filtering procedure for eddy-resolving simulations with a spectral element ocean model, *J. Comput. Phys.* **137**(1), 130–154.

Lewis, D. L., Lund, J. and Bowers, K. L.: 1987, The space-time sinc-Galerkin method for parabolic problems, *International Journal for Numerical Methods in Engineering* **24**, 1629–1644. Sinc basis in *both* space and time.

Lewis, H. R. and Bellan, P. M.: 1990, Physical constraints on the coefficients of Fourier expansions in cylindrical coordinates, *Journal of Mathematical Physics* **31**, 2592–. Analysis of the symmetry conditions across $r = 0$ for both scalars and vectors.

Li, C. W. and Qin, M. Z.: 1988, A symplectic difference scheme for the infinite dimensional Hamiltonian system, *Journal of Computer and Applied Mathematics* **6**, 164–174.

Li, T.-Y.: 1987, Solving polynomial systems, *Math. Intelligencer* **9**, 33–39. Probability-one homotopy.

Lie, I.: 1993, Using implicit ODE methods with iterative linear equation solvers in spectral methods, *SIAM Journal of Scientific Computing* **14**, 1194–1213.

Liffman, K.: 1996, Comments on a collocation spectral solver for the Helmholtz equation, *Journal of Computational Physics* **128**(1), 254–258. Extends the linear, elliptic boundary value solver of Ehrenstein and Peyret(1986), which uses Chebyshev polynomials, to inhomogeneous Robin boundary conditions.

Lin, S.-H. and Pierrehumbert, R. T.: 1988, Does Ekman friction suppress baroclinic instability?, *Journal of the Atmospheric Sciences* **45**, 2920–2933. Solves a two-dimensional eigenvalue problem on the half-space ($[-\infty, \infty] \otimes [0, \infty]$) via a tensor product basis of rational Chebyshev functions, $TB_m(y)\,TL_n(z)$.

Lindberg, C. and Broccoli, A. J.: 1996, Representation of topography in spectral climate models and its effect on simulated precipitation, *J. Climate* **9**, 2641–. Smoothing of spherical harmonic series for topography through non-uniform spherical spline.

Lindzen, R. S.: 1970, Internal equatorial planetary-scale waves in shear flow, *Journal of the Atmospheric Sciences* **27**, 394–407. Mixed elliptic-hyperbolic PDE.

Liu, K. M. and Ortiz, E. L.: 1982, Eigenvalue problems for singularly perturbed differential equations, *in* J. J. H. Miller (ed.), *Computational Methods for Boundary and Interior Layers, Vol. II*, Boole Press, Dublin, pp. 324–329.

Liu, K. M. and Ortiz, E. L.: 1986, Numerical solution of eigenvalue problems for partial differential equations with tau-lines method, *Computer Mathematics and Applications* **12B**, 1153–1168.

Liu, K. M. and Ortiz, E. L.: 1987a, Tau method approximate solution of high-order differential eigenvalue problems defined in the complex plane with an application of the Orr-Sommerfeld stability equation, *Comm. Appl. Numer. Meths.* **3**, 187–194.

Liu, K. M. and Ortiz, E. L.: 1987b, Tau method approximation of differential eigenvalue problems where the spectral parameter enters nonlinearly, *Journal of Computational Physics* **72**(2), 299–310.

Liu, K. M. and Ortiz, E. L.: 1989, Numerical solution of ordinary and partial functional-differential eigenvalue problems with the tau method, *Computing* **41**, 205–217.

Liu, K. M., Ortiz, E. L. and Pun, K.-S.: 1984, Numerical solution of Steklov's partial differential equation eigenvalue problem, *in* J. J. H. Miller (ed.), *Computational Methods for Boundary and Interior Layers, Vol. II*, Boole Press, Dublin, pp. 244–249.

Liu, Y., Liu, L. and Tang, T.: 1994, The numerical computation of connecting orbits in dynamical systems: A rational spectral approach, *Journal of Computational Physics* **111**, 373–380. TB_n basis for computing heteroclinic and homoclinic solutions (i. e., shock-like and soliton-like) solutions to nonlinear boundary value problems on an infinite interval.

Lomtev, I., Quillen, C. B. and Karniadakis, G. E.: 1998, Spectral/hp methods for viscous compressible flows on unstructured 2D meshes, *J. Comput. Phys.* **144**(2), 325–357. Spectral elements on triangles using a discontinuous Galerkin formulation.

Longuet-Higgins, M. S.: 1968, The eigenfunctions of Laplace's tidal equation over a sphere, *Phil. Trans. Royal Society of London, Series A* **262**, 511–607. Spherical harmonics; tridiagonal Galerkin matrices.

Lopez, J. M. and Shen, J.: 1998, An efficient spectral-projection method for the Navier-Stokes equations in cylindrical coordinates, *Journal of Computational Physics* **139**, 308–326.

Lorenz, E. N. and Krishnamurthy, V.: 1987, On the nonexistence of a slow manifold, *Journal of the Atmospheric Sciences* **44**, 2940–2950. Weakly non-local in time; nothing on spectral methods except that the five-mode system is derived by a Galerkin method.

Lund, J.: 1986, Symmetrization of the sinc-Galerkin method for boundary value problems, *Mathematics of Computation* **47**, 571–588.

Lund, J. and Bowers, K. L.: 1992, *Sinc Methods For Quadrature and Differential Equations*, Society for Industrial and Applied Mathematics, Philadelphia. 304 pp. Restricted to sinc functions only.

Lund, J. and Vogel, C. R.: 1990, A fully-Galerkin method for the numerical solution of an inverse problem in a parabolic partial differential equation, *Inverse Problems* **6**, 205–217. Sinc basis.

Lund, J., Bowers, K. L. and Carlson, T. S.: 1991, Fully sinc-Galerkin computation for boundary feedback stabilization, *Journal of Mathematical Systems, Estimation and Control* **1**, 165–182.

Lund, J., Bowers, K. L. and McArthur, K. M.: 1989, Symmetrization of the sinc-Galerkin method with block techniques for elliptic equations, *IMA Journal of Numerical Analysis* **9**, 29–46.

Lund, J. R. and Riley, B. B.: 1984, A sinc-collocation method for the computation of the eigenvalues of the radial Schroedinger equation, *IMA Journal of Numerical Analysis* **4**, 83–98.

Lundin, L.: 1980, A cardinal function method of solution of the equation $\Delta u = u - u^3$, *Mathematics of Computation* **35**, 747–756. Sinc (Whittaker cardinal) basis.

Lynch, P.: 1992, Richardson's barotropic forecast: A reappraisal, *Bulletin of the American Meteorological Society* **73**(1), 35–47.

Lynch, P. and Huang, X.: 1992, Initialization of the HIRLAM model using a digital filter, *Monthly Weather Review* **120**, 1019–1034. Not spectral; slow manifold initialization scheme.

Ma, H.: 1992, The equatorial basin response to a Rossby wave packet: The effects of nonlinear mechanism, *Journal of Marine Research* **50**, 567–609. Spectral element ocean model.

Ma, H.: 1993a, Trapped internal gravity waves in a geostrophic boundary current, *Journal of Fluid Mechanics* **247**, 205–229. Spectral element ocean model.

Ma, H.: 1993b, A spectral element basin model for the shallow water equations, *Journal of Computational Physics* **109**, 133–149. Spectral element ocean model.

Ma, H.: 1995, Parallel computation with the spectral element method, *in* A. Ecer, J. Periaux, N. Satofuka and S. Taylor (eds), *Parallel Computational Fluid Dynamics: Implementations and Results Using Parallel Computers*, Elsevier, Amsterdam, pp. 239–246.

Ma, H.: 1996a, Baroclinic wave motions in an equatorial ocean current and related temperature fluctuations, *J. Marine Res.* **54**(6), 1073–1096. Spectral elements.

Ma, H.: 1996b, The dynamics of North Brazil Current retroflection eddies, *Journal of Marine Research* **54**, 35–53. Spectral elements.

Ma, H.-P. and Guo, B.-Y.: 1986, The Fourier pseudo-spectral method with a restrain operator for the Korteweg-de Vries equation, *Journal of Computational Physics* **65**, 120–137.

Ma, H.-P. and Guo, B.-Y.: 1987, The Fourier pseudo-spectral method for solving two-dimensional vorticity equations, *IMA Journal of Numerical Analysis* **5**, 47–60.

Machenhauer, B.: 1977, On the dynamics of gravity oscillations in a shallow water model, with application to normal mode initialization, *Beitr. Phys. Atmos.* **50**, 253–271.

Machenhauer, B. and Daley, R.: 1972, A baroclinic primitive equation model with a spectral representation in three dimensions, *Report 4*, Institut for Teoretisch Meteorologi, Copenhagen University, Copenhagen, Denmark. 66 pp. Legendre series in the vertical, spherical harmonics in latitude and longitude.

Machenhauer, B. and Daley, R.: 1974, Hemispheric spectral model, number 14 in *GARP Publication Series*, World Meteorological Organization, Geneva, Switzerland, pp. 226–251. Weather forecasting model with Legendre polynomials in height and spherical harmonics in latitude and longitude.

Maday, Y. and Patera, A. T.: 1987, Spectral element methods for the incompressible Navier Stokes equations, *in* A. K. Noor and J. T. Oden (eds), *State of the Art Surveys on Computational Mechanics*, ASME, New York, pp. 71–143.

Maday, Y. and Rønquist, E. M.: 1990, Optimal error analysis of spectral methods with emphasis on non-constant coefficients and deformed geometries, *in* C. Canuto and A. Quarteroni (eds), *Spectral and High Order Methods for Partial Differential Equations: Proceedings of the ICOSAHOM '89 Conference in Como, Italy*, North-Holland/Elsevier, Amsterdam, pp. 91–115. Also in Comput. Meths. Appl. Mech. Engrg., vol. 80, with the same page numbers.

Maday, Y., Mavriplis, C. A. and Patera, A. T.: 1989, Nonconforming mortar element method: application to spectral discretizations, *in* T. F. Chan, R. Glowinski, J. Periaux and O. B. Widlund (eds), *Domain Decomposition Methods*, SIAM, Society for Industrial and Applied Mathematics SIAM, Philadelphia.

Maday, Y., Meiron, D., Patera, A. T. and Rønquist, E. M.: 1993, Analysis of iterative methods for the steady and unsteady Stokes problem: Application to spectral element discretizations, *SIAM Journal of Scientific Computing* **14**, 310–337.

Maday, Y., Patera, A. T. and Rønquist, E. M.: 1990, An operator integration-factor splitting method for time-dependent problems: Application to incompressible fluid flow, *Journal of Scientific Computing* **5**, 263–292. Generalizes the usual two-level and three-level schemes, which are of at most second order accuracy in time, to arbitrary order.

Maday, Y., Patera, A. T. and Rønquist, E. M.: 1992, The $P_N \times P_{N-2}$ method for the approximation of the Stokes problem, *Laboratoire d'Analyse Numérique, Paris VI* **11**, 4.

Maday, Y., Pernaud-Thomas, B. and Vandeven, H.: 1985, Reappraisal of Laguerre type spectral methods, *Rech. Aerosp.* **1985-6**, 13–35.

Makar, P. A. and Karpik, S. R.: 1996, Basis-spline interpolation on the sphere: application to semi-Lagrangian advection, *Monthly Weather Review* **124**, 182. Comparisons with spectral methods.

Malek, A. and Phillips, T. N.: 1995, Pseudospectral collocation methods for fourth-order differential equations, *IMA Journal of Numerical Analysis* **15**, 523–553.

Malik, M. R.: 1990, Numerical methods for hypersonic boundary layer stability, *Journal of Computational Physics* **86**(2), 376–413. Linear eigenvalue problems solved by single domain and multi-domain Chebyshev methods and compact 4th order differences; inverse Rayleigh iteration.

Malik, M. R. and Orszag, S. A.: 1987, Linear stability analysis of three-dimensional compressible boundary layers, *Journal of Scientific Computing* **2**, 77–98.

Mamun, C. K. and Tuckerman, L. S.: 1995, Asymmetry and Hopf bifurcation in spherical Couette flow, *Physics of Fluids* **7**(1), 80–91. Axisymmetric incompressible flow between two spheres using vorticity/streamfunction and influence matrix method. Good discussion of how, by modifying only twenty lines in the time-dependent code, steady states could be computed directly by a nonlinear Richardson's iteration with preconditioning by Stokes flow (that is, by inverting the operator which is treated implicitly in the semi-implicit time-marching algorithm).

Mansell, G., Merryfield, W., Shizgal, B. and Weinert, U.: 1993, A comparison of differential quadrature methods for the solution of partial differential equations, *Computer Methods in Applied Mechanics and Engineering* **104**, 295–316. Differential quadrature is the same or almost the same as the pseudospectral method.

Marchuk, G. I.: 1974, *Numerical Methods in Numerical Weather Prediction*, Academic Press, New York. 277 pp. Splitting and fractional steps time integration.

Marcus, P. A.: 1984a, Simulation of Taylor-Couette flow, Part 1. Numerical methods and comparison with experiment, *Journal of Fluid Mechanics* **146**, 45–64. Fourier-Fourier-Chebyshev computations of flow in an annulus in cylindrical coordinates, assumed periodic in z as well as θ. "Shift-and-reflect" symmetry halves the number of Fourier modes in the basis.

Marcus, P. A.: 1984b, Simulation of Taylor-Couette flow, Part 2. Numerical results for wavy vortex flow with one travelling wave, *Journal of Fluid Mechanics* **146**, 65–113.

Marcus, P. A. and Tuckerman, L. S.: 1987a, Simulation of flow between concentric rotating spheres. Part 1. Steady states, *J. Fluid Mech.* **185**, 1–30. Axisymmetric flows with splitting errors removed by a Green's function [influence matrix] method.

Marcus, P. A. and Tuckerman, L. S.: 1987b, Simulation of flow between concentric rotating spheres. Part 2. Transitions, *J. Fluid Mech.* **185**, 31–65.

Marcus, P. A., Orszag, S. A. and Patera, A. T.: 1983, Simulation of cylindrical Couette flow, *in* E. Krause (ed.), *8th International Conf. on Numerical Methods in Fluid Dynamics*, number 170 in *Lecture Notes in Physics*, Springer-Verlag, New York, pp. 371–376.

Marcus, P. J.: 1990, Vortex dynamics in a shearing zonal flow, *Journal of Fluid Mechanics* **215**, 393–430. Fourier-Chebyshev algorithm for annular flow from Marcus(1984a). Dealiasing using Orszag Two-Thirds Rule because it helped here, contrary to the experiments of Marcus(1984b). Compared hyperviscosity with filtering of high order modes, and preferred the latter because it was cheaper, not requiring additional boundary conditions.

Margolin, L. G. and Jones, D. A.: 1992, An approximate inertial manifold for computing Burgers' equation, *Physica D* **60**, 175–184.

Marion, M. and Témam, R.: 19, Nonlinear Galerkin methods, *SIAM Journal of Numerical Analysis* **26**, 1139–1157.

Marshall, H. G. and Boyd, J. P.: 1987, Solitons in a continuously stratified equatorial ocean, *Journal of Physical Oceanography* **17**, 1016–1031. Hermite function application.

Mason, J. C.: 1967, Chebyshev polynomial approximations for the L-shaped membrane eigenvalue problem, *SIAM Journal of Applied Mathematics* **15**, 172–186. Maps the L-shaped domain into a polygonal figure and then solves the problem on the square containing the polygon. This completely eliminates the corner singularities; his results are superb.

Matsushima, T. and Marcus, P. S.: 1995, A spectral method for polar coordinates, *Journal of Computational Physics* **120**, 365–374. One-side Jacobi polynomials in radius.

Matsushima, T. and Marcus, P. S.: 1997, A spectral method for unbounded domains, *Journal of Computational Physics* **137**(2), 321–345. Polar coordinates with Fourier basis in angle and a radial basis of rational functions which are the images of associated Legendre functions under an algebraic mapping; applications to hydrodynamic vortices.

Mavriplis, C.: 1989, Laguerre polynomials for infinite-domain spectral elements, *Journal of Computational Physics* **80**(2), 480–.

Mavriplis, C.: 1994, Adaptive mesh strategies for the spectral element method, *in* C. Bernardi and Y. Maday (eds), *Analysis, Algorithms and Applications of Spectral and High Order Methods for Partial Differential Equations*, Selected Papers from the International Conference on Spectral and High Order Methods (ICOSAHOM '92), Le Corum, Montpellier, France, 22-26 June 1992, North-Holland, Amsterdam, pp. 77–86. Also in Comput. Meths. Appl. Mech. Engr., vol. 116.

Mayer, E. W. and Powell, K. G.: 1992, Viscous and inviscid instabilities of a trailing vortex, *Journal of Fluid Mechanics* **245**, 91–114. Chebyshev polynomial/QR one-dimensional eigenvalue problem in radius r in polar coordinates with domain truncation at large but finite r. Integration on a contour deformed into the complex plane is used to resolve nearly-neutral modes. The infinite domain is truncated to an annulus for some runs to calculate "ring modes" which are concentrated at intermediate radius.

McArthur, K. M., Bowers, K. L. and Lund, J.: 1987a, Numerical implementation of the sinc-Galerkin method for second-order hyperbolic equations, *Numerical Methods for Partial Differential Equations* **3**, 169–185.

McArthur, K. M., Bowers, K. L. and Lund, J.: 1987b, The sinc method in multiple space dimensions: Model problems, *Numerische Mathematik* **56**, 789–816.

McCalpin, J. D.: 1988, A quantitative analysis of the dissipation inherent in Semi-Lagrangian advection, *Monthly Weather Review* **116**, 2330–2336. Not spectral, but good analysis of the damping which is implicit in off-grid interpolation.

McCrory, R. L. and Orszag, S. A.: 1980, Spectral methods for multi-dimensional diffusion problems, *Journal of Computational Physics* **37**, 93–112.

McDonald, A.: 1984, Accuracy of multiply-upstream semi-Lagrangian advective schemes I., *Monthly Weather Review* **112**, 1267–1275. Not spectral.

McDonald, A.: 1986, A semi-Lagrangian and semi-implicit two-time-level integration scheme, *Monthly Weather Review* **114**, 824–830. Not spectral; only first order accurate in time.

McDonald, A.: 1987, Accuracy of multiply-upstream semi-Lagrangian advection schemes II, *Monthly Weather Review* **115**, 1446–1450. Not spectral.

McDonald, A. and Bates, J. R.: 1987, Improving the estimate of the departure point position in a two-time-level semi-Lagrangian and semi-implicit model, *Monthly Weather Review* **115**, 737–739. Not spectral.

McDonald, A. and Bates, J. R.: 1989, Semi-Lagrangian integration of a gridpoint shallow-water model on the sphere, *Monthly Weather Review* **117**(1), 130–137. Not spectral.

McFadden, G. B., Murray, B. T. and Boisvert, R. F.: 1990, Elimination of spurious eigenvalues in the Chebyshev tau spectral method, *Journal of Computational Physics* **91**, 228–239.

McGregor, J. L.: 1993, Economical determination of departure points for semi-Lagrangian models, *Monthly Weather Review* **121**, 221–230. Not spectral per se; shows his $O(\tau^2)$ formula D_2 costs only 5 additions and 7 multiplications versus 30 and 60 for bicubic interpolation (in two space dimensions).

McKerrell, A.: 1988, The global element applied to fluid flow problems, *Computers and Fluids* **16**, 41–46. Corner singularities, domain decomposition algorithms, two-dimensional mappings.

McKerrell, A. and Delves, L. M.: 1984, Solution of the global element equations on the ICL DAP, *ICL Technical Journal* **4**, 50–58. Massively parallel computation with spectral domain decomposition.

McKerrell, A., Phillips, C. and Delves, L. M.: 1981, Chebyshev expansion methods for the solution of elliptic partial differential equations, *Journal of Computational Physics* **40**, 444–452.

McLaughlin, J. B. and Orszag, S. A.: 1982, Transition from periodic to chaotic thermal convection, *Journal of Fluid Mechanics* **122**, 123–142.

Mead, K. O. and Delves, L. M.: 1973, On the convergence rate of generalized Fourier expansions, *Journal of the Institute for Mathematics and Its Applicaitons* **12**, 247–259.

Meiron, D. I., Baker, G. R. and Orszag, S. A.: 1982, Analytic structure of vortex sheet dynamics. Part 1. Kelvin-Helmholtz instability, *J. Fluid Mech.* **114**, 283–298. Singularities estimated from the slopes of the Fourier coefficients versus degree at different times.

Meiron, D. I., Orszag, S. A. and Israeli, M.: 1981, Applications of numerical conformal mapping, *Journal of Computational Physics* **40**, 345–360.

Mercier, B.: 1989, *An Introduction to the Numerical Analysis of Spectral Methods*, Vol. 318 of *Lecture Notes in Physics*, Springer-Verlag, New York. 200 pp. Author's translation, without updating, of an earlier monograph in French.

Mercier, B. and Raugel, G.: 1982, Resolution d'un problème aux limites dans un ouvert axisymétrique par éléments finis en r, z et séries de Fourier en théta, *R. A. I. R. O. Anal. Numer.* **16**, 67–100.

Merilees, P. E.: 1973a, An alternative scheme for the summation of a series of spherical harmonics, *J. Appl. Meteor.* **12**, 224–227. Instability of standard recurrence for the spherical harmonics for large zonal wavenumber; ill-conditioning of "Robert" basis.

Merilees, P. E.: 1973b, The pseudospectral approximation applied to the shallow water wave equations on a sphere, *Atmosphere* **11**, 13–20.

Merilees, P. E.: 1974, Numerical experiments with the pseudospectral method in spherical coordinates, *Atmosphere* **12**, 77–96.

Merzbacher, E.: 1970, *Quantum Mechanics*, 2 edn, Wiley, New York. 400 pp. Hermite funcs.; Galerkin meths. in quantum mechanics.

Metcalfe, R. W., Orszag, S. A., Brachet, M. E., Menon, S. and Riley, J.: 1987, Secondary instability of a temporally growing mixing layer, *Journal of Fluid Mechanics* **184**, 207–243.

Mills, R. D.: 1987, Using a small algebraic manipulation system to solve differential and integral equations by variational and approximation techniques, *J. Symbolic Comp.* **3**, 291–301.

Minion, M. L. and Brown, D. L.: 1997, Performance of under-resolved two-dimensional incompressible flow simulations, II, *J. Comput. Phys.* **138**(2), 734–765. Comparison of several numerical methods including a pseudospectral method; generation of spurious vortices whose size is much larger than twice the grid spacing through instability, excited by truncation error, of underresolved flows.

Mittal, R. and Balachandar, S.: 1996, Direct numerical simulation of flow past elliptic cylinders, *Journal of Computational Physics* **124**, 351–367. Chebyshev-Fourier time-dependent calculations using a single computational domain in elliptic coordinates. Good, careful discussion of both inflow and outflow boundary conditions, parabolization of the outflow, smooth blending of outflow into inflow.

Mizzi, A., Tribbia, J. and Curry, J.: 1995, Vertical spectral representation in primitive equation models of the atmosphere, *Monthly Weather Review* **123**(8), 2426–2446. Removed most obstacles to use of spectral methods in z, but their series of vertical normal modes converged rather slowly.

Moore, D. W. and Philander, S. G. H.: 1977, Modelling of the tropical oceanic circulation, *in* E. D. Goldberg (ed.), *The Sea, Volume 6*, Wiley, New York, pp. 319–361. Hermite functions.

Moore, S., Healy, Jr, D. M. and Rockmore, D.: 1993, Symmetry stabilization for fast discrete monomial transforms and polynomial evaluations, *Lin. Alg. Appl.* **192**, 249–299. $O(N \log_2(N))$ discrete Fourier transform on nonuniform grids.

Morf, R., Orszag, S. A. and Frisch, U.: 1980, Spontaneous singularity in three-dimensional, inviscid compressible flow, *Phys. Rev. Lett.* **44**, 572–575. The conclusion that singularities can form was (mostly) repudiated in Brachet *et al.*(1983).

Morgan, A. P.: 1987, *Solving Polynomial Systems Using Continuation for Scientific and Engineering Problems*, Prentice-Hall, Englewood Cliffs, New Jersey. Not spectral.

Morse, P. M. and Feshbach, H.: 1953, *Methods of Theoretical Physics*, McGraw-Hill, New York. 2000 pp, (in two volumes) Good treatise on solving linear partial differential equations through the method of separation of variables.

Morton, K. W.: 1985, Generalised Galerkin methods for hyperbolic problems, *Computer Methods in Applied Mechanics and Engineering* **52**, 847–871. Not spectral; independent invention of a semi-Lagrangian method.

Mulholland, L. and Sloan, D.: 1991, The effect of filtering on the pseudospectral solution of evolutionary partial differential equations, *Journal of Computational Physics* **96**, 369–390.

Mulholland, L. S. and Sloan, D. M.: 1992, The role of preconditioning in the solution of evolutionary partial differential equations by implicit Fourier pseudospectral methods, *Journal of Computational and Applied Mathematics* **42**, 157–174.

Mulholland, L. S., Huang, W.-Z. and Sloan, D. M.: 1998, Pseudospectral solution of near-singular problems using numerical coordinate transformations based on adaptivity, *SIAM J. Sci. Comput.* **19**(4), 1261–1289.

Mullholland, L. S., Qiu, Y. and Sloan, D. M.: 1997, Solution of evolutionary PDEs using adaptive finite differences with pseudospectral post-processing, *J. Comput. Phys.* **131**(2), 280–298.

Nakamura, S.: 1996, *Numerical Analysis and Graphic Visualization with MATLAB*, Prentice-Hall, Upper Saddle River, New Jersey. Not spectral.

Namasivayam, S. and Ortiz, E. L.: 1981, Best approximation and the numerical solution of partial differential equations with the Tau method, *Portug. Math.* **40**, 97–119.

Namasivayam, S. and Ortiz, E. L.: 1993, Error analysis of the Tau method: dependence of the approximation error on the choice of perturbation term, *Computers Math. applic.* **25**(1), 89–104.

Navarra, A.: 1987, An application of the Arnoldi's method to a geophysical fluid dynamics problem, *Journal of Computational Physics* **69**, 143–162. Mixed finite difference/spherical harmonics Galerkin treatment of boundary value problem; Arnoldi's method allows computations of unstable modes with as many as 13,000 unknowns.

Navarra, A., Stern, W. F. and Miyakoda, K.: 1994, Reduction of the Gibbs oscillation in spectral model simulations, *J. Climate* **7**, 1169–1183. Apply several different smoothers to spherical harmonics series with sucess. Note that the usual second order finite differences can be equivalently derived by applying Lanczos smoothing to the spectral sum for a derivative.

Navon, I. M.: 1987, PENT: A periodic pentadiagonal systems solver, *Comm. Appl. Numer. Meths.* **3**, 63–69. Not spectral, but useful matrix-solver.

Nicholls, D. P.: 1998, Travelling water waves: Spectral continuation methods with parallel implementation, *J. Comput. Phys.* **143**(1), 224–240. Fourier pseudospectral solution to a nonlinear eigenvalue problem.

Nosenchuck, D. M., Krist, S. E. and Zang, T. A.: 1987, Multigrid methods for the Navier-Stokes Computer, *in* S. McCormick and K. Stuben (eds), *Multigrid Methods*, Marcel Dekker, New York, pp. 491–516.

Nouri, F. Z. and Sloan, D. M.: 1989, A comparison of Fourier pseudospectral methods for the solution of the Korteweg-deVries equation, *Journal of Computational Physics* **83**, 324–344.

O'Connor, W. P.: 1995, The complex wavenumber eigenvalues of Laplace's tidal equations for oceans bounded by meridians, *Proceedings of the Royal Society of London A* **449**, 51–64. Associated Legendre functions through a recurrence-derived Galerkin methods, which is susceptible to errors [as confirmed by Corrigendum: O'Connor(1996).].

O'Connor, W. P.: 1996, The complex wavenumber eigenvalues of Laplace's tidal equations for oceans bounded by meridians: Corrigendum, *Proceedings of the Royal Society of London A* **452**, 1185–1187.

Orszag, S. A.: 1970, Transform method for calculation of vector coupled sums: Application to the spectral form of the vorticity equation, *Journal of the Atmospheric Sciences* **27**, 890–895. Implementing spectral methods in spherical harmonics.

Orszag, S. A.: 1971a, On the elimination of aliasing in finite difference schemes by filtering high-wavenumber components, *Journal of the Atmospheric Sciences* **28**, 1074. A two-paragraph classic.

Orszag, S. A.: 1971b, Accurate solution of the Orr-Sommerfeld equation, *Journal of Fluid Mechanics* **50**, 689–703. Combines Chebyshev method with QR or QZ matrix eigen-solver to solve a linear stability problem.

Orszag, S. A.: 1971c, Numerical simulations of incompressible flows within simple boundaries: accuracy, *Journal of Fluid Mechanics* **49**, 75–112.

Orszag, S. A.: 1971d, Numerical simulations of incompressible flows within simple boundaries: Galerkin (spectral) representations, *Studies in Applied Mathematics* **50**, 293–327.

Orszag, S. A.: 1971e, Galerkin approximations to flows within slabs, spheres and cylinders, *Physical Review Letters* **26**, 1100–1103.

Orszag, S. A.: 1972, Comparison of pseudospectral and spectral approximations, *Studies in Applied Mathematics* **51**, 253–259.

Orszag, S. A.: 1974, Fourier series on spheres, *Monthly Weather Review* **102**, 56–75.

Orszag, S. A.: 1976, Turbulence and transition: A progress report, *in* A. I. vander Vooren and Zandbergen (eds), *Proceedings of the Fifth International Conference on Numerical Fluid Dynamics*, number 59 in *Lecture Notes in Physics*, Springer-Verlag, New York, pp. 39–51.

Orszag, S. A.: 1979, Spectral methods for problems in complex geometries, *in* S. V. Parter (ed.), *Numerical Methods for Partial Differential Equations*, Academic Press, New York. Shorter, preliminary version of Orszag (1980).

Orszag, S. A.: 1980, Spectral methods for problems in complex geometries, *Journal of Computational Physics* **37**, 70–92. Independent invention of finite difference preconditioning for solving boundary value problems by means of Chebyshev polynomials and iteration.

Orszag, S. A.: 1986, Fast eigenfunction transforms, *in* G. C. Rota (ed.), *Science and Computers*, Academic Press, New York, pp. 23–30. A generalization of the Fast Fourier Transform to functions that satisfy three-term recurrence relations.

Orszag, S. A. and Israeli, M.: 1974, Numerical simulation of incompressible flow, *Ann. Revs. Fluid Mech.* **6**, 281–318. REVIEW.

Orszag, S. A. and Kells, L. C.: 1980, Transition to turbulence in plane Poiseuille flow and plane Couette flow, *Journal of Fluid Mechanics* **96**, 159–205.

Orszag, S. A. and Patera, A. T.: 1980, Subcritical transition to turbulence in plane channel flows, *Physical Review Letters* **45**, 989–993.

Orszag, S. A. and Patera, A. T.: 1981, Subcritical transition to turbulence in planar shear flow, *in* R. E. Meyer (ed.), *Transition and Turbulence*, Academic Press, New York, pp. 127–146.

Orszag, S. A. and Patera, A. T.: 1983, Secondary instability of wall bounded shear flows, *Journal of Fluid Mechanics* **128**, 347–385.

Orszag, S. A. and Paterson, Jr., G. S.: 1972a, *Statistical Models of Turbulence*, Springer-Verlag, New York, chapter Numerical simulation of turbulence, pp. 127–147.

Orszag, S. A. and Paterson, Jr., G. S.: 1972b, Numerical simulation of three dimensional homogeneous isotropic turbulence, *Phys. Rev. Lett.* **28**, 76–79.

Orszag, S. A. and Tang, C. M.: 1979, Small-scale structure of two-dimensional magnetohydrodynamic turbulence, *Journal of Fluid Mechanics* **90**, 129–143.

Orszag, S. A., Israeli, M. and Deville, M.: 1986, Boundary conditions for incompressible flows, *Journal of Scientific Computing* **1**, 75–111. Thorough discussion of high order artificial boundary conditions for fractional step (splitting) time-marching so that the computed flows are both nondivergent and no-slip.

Ortiz, E. L.: 1969, The tau method, *SIAM Journal of Numerical Analysis* **6**, 480–492.

Ortiz, E. L.: 1987, Recent progress in the numerical treatment of singular problems for partial differential equations with techniques based on the Tau method, *in* E. L. Ortiz (ed.), *Numerical Approximations of P. D. E., Part III*, North-Holland, Amsterdam, pp. 83–98.

Ortiz, E. L. and Pham Ngoc Dinh, A.: 1987, Linear recursive schemes associated with some nonlinear partial differential equations in one dimension and the tau method, *SIAM J. Math. Anal.* **18**(2), 452–464.

Ortiz, E. L. and Pun, K.-S.: 1985, Numerical solution of nonlinear partial differential equations with the tau method, *Journal of Computational and Applied Mathematics* **12 and 13**, 511–516.

Ortiz, E. L. and Samara, H.: 1981, An operational approach to the Tau method for the numerical solution of nonlinear differential equations, *Computing (Wien)* **27**, 15–25.

Ortiz, E. L. and Samara, H.: 1983, Numerical solution of differential eigenvalue problems with an operational approach to the tau method, *Computing (Wien)* **31**, 95–103.

Owens, R. G.: 1998, Spectral approximations on the triangle, *Proc. R. Soc. Lond. A* **454**, 857–872. Derives a new polynomial basis for the triangle through a Sturm-Liouville eigenproblem in two dimensions; the eigenfunctions are products of Legendre polynomials in one transformed coordinate with hypergeometric polynomials in the other. A spectral rate of convergence is proved. The polynomials are used to generate a new cubature formula for the triangle.

Owens, R. G. and Phillips, T. N.: 1991, A spectral domain decomposition method for the planar non-Newtonian stick-slip problem, *Journal of Non-Newtonian Fluid Mechanics* **41**, 43–79.

Owens, R. G. and Phillips, T. N.: 1996, Steady viscoelastic flow past a sphere using spectral elements, *International Journal of Numerical Methods in Engineering* **39**(9), 1517–1534. Flow past sphere embedded in a cylindrical tube.

Panchang, V. G. and Kopriva, D. A.: 1989, Solution of two-dimensional water-wave propagation problems by Chebyshev collocation, *Mathematical and Computer Modelling* **12**, 625–640.

Pascal, F. and Basdevant, C.: 1992, Nonlinear Galerkin method and subgrid-scale model for two-dimensional turbulent flows, *Theoretical and Computational Fluid Dynamics* **3**, 267–284.

Patera, A. T.: 1984, A spectral element method for fluid dynamics: laminar flow in a channel expansion, *Journal of Computational Physics* **54**, 468–488.

Patera, A. T.: 1986, Fast direct Poisson solvers for high-order finite element discretizations in rectangularly decomposable domains, *Journal of Computational Physics* **65**, 474–480. Combines the eigenfunction scheme of Haidvogel and Zang(1979) with static condensation to create a direct method of $O(N^{5/2})$ cost.

Patera, A. T. and Orszag, S. A.: 1980, Transition and turbulence in plane channel flows, *in* R. W. MacCormack and W. C. Reynolds (eds), *Proceedings of the 7th International Conference on Numerical Methods in Fluid Dynamics*, Springer-Verlag, New York, pp. 329–335.

Patera, A. T. and Orszag, S. A.: 1981, Finite-amplitude stability of axisymmetric pipe flow, *Journal of Fluid Mechanics* **112**, 467.

Patera, A. T. and Orszag, S. A.: 1986, Instability of pipe flow, *in* A. R. Bishop, D. K. Campbell and B. Nicolaenko (eds), *Nonlinear Problems: Present and Future*, North Holland, Amsterdam, pp. 367–377.

Patterson Jr., G. S. and Orszag, S. A.: 1971, Spectral calculation of isotropic turbulence: Efficient removal of aliasing interaction, *Physics of Fluids* **14**, 2538–2541.

Pellerin, P., Laprise, R. and Zawadzki, I.: 1995, The performance of a semi-Lagrangian transport scheme for the advection-condensation problem, *Monthly Weather Review* **123**, 3318.

Pelz, R. B.: 1993, Parallel compact FFTs for real sequences, *SIAM J. Sci.Comput.* **14**(4), 914–935.

Phillips, N. J.: 1956, The general circulation of the atmosphere: A numerical experiment, *Quart. J. Roy. Met. Soc.* **82**, 123–164. The empirical discovery of aliasing instability.

Phillips, N. J.: 1959, An example of nonlinear computational instability, *The Atmosphere and the Sea in Motion*, Rockefeller Institute Press, New York, pp. 501–504. The theoretical discovery of aliasing instability.

Phillips, T. N.: 1988, On the Legendre coefficients of a general-order derivative of an infinitely differentiable function, *IMA Journal of Numerical Analysis* **8**, 455–459.

Phillips, T. N. and Davies, A. R.: 1988, On semi-infinite spectral elements for poisson problems with re-entrant boundary singularities, *Journal of Computational and Applied Mathematics* **21**, 173–188.

Phillips, T. N. and Karageorghis, A.: 1990, On the coefficients of integrated expansions of ultraspherical polynomials, *SIAM Journal on Numerical Analysis* **27**, 823–830.

Phillips, T. N. and Owens, R. G.: 1997, A mass conserving multi-domain spectral collocation method for the Stokes problem, *Comput. Fluids* **26**(8), 825–840.

Phillips, T. N., Zang, T. A. and Hussaini, M. Y.: 1986, Preconditioners for the spectral multigrid method, *IMA Journal of Numerical Analysis* **6**, 273–292.

Pinelli, A., Benocci, C. and Deville, M.: 1992, A preconditioing technique for Chebyshev collocated advection diffusion operators, *in* C. Hirsh (ed.), *First European CFD Conference Proceedings*, Elsevier, Amsterdam, p. 109.

Pinelli, A., Benocci, C. and Deville, M.: 1994, A Chebyshev collocation algorithm for the solution of advection-diffusion equations, *in* C. Bernardi and Y. Maday (eds), *Analysis, Algorithms and Applications of Spectral and High Order Methods for Partial Differential Equations*, Selected Papers from the International Conference on Spectral and High Order Methods (ICOSAHOM '92), Le Corum, Montpellier, France, 22-26 June 1992, North-Holland, Amsterdam, pp. 201–210. Also in Comput. Meths. Appl. Mech. Engrg., vol. 116.

Pinelli, A., Couzy, W., Deville, M. O. and Benocci, C.: 1996, An efficient iterative solution method for the Chebyshev collocation of advection-dominated transport problems, *SIAM Journal of Scientific Computing* **17**(3), 647–657.

Pironneau, O.: 1982, On the transport-diffusion algorithm and its application to the Navier-Stokes equations, *Numerische Mathematik* **38**, 309–322. Not spectral; invention of a semi-Lagrangian method.

Press, W. H., Flannery, B. H., Teukolsky, S. A. and Vetterling, W. T.: 1986, *Numerical Recipes: The Art of Scientific Computing*, Cambridge University Press, New York.

Priestley, A.: 1993, A quasi-conservative version of the semi-Lagrangian advection scheme, *Monthly Weather Review* **121**, 621. Not spectral.

Priymak, V. G.: 1995, Pseudospectral algorithms for Navier-Stokes simulation of turbulent flows in cylindrical geometry with coordinate singularities, *Journal of Computational Physics* **118**, 366–379.

Priymak, V. G. and Miyazaki, T.: 1998, Accurate Navier-Stokes investigation of transitional and turbulent flows in a circular pipe, *Journal of Computational Physics* **142**, 370–411. Chebyshev in radius, Fourier in the other two coordinates.

Promislow, K. and Témam, R.: 1991, Approximate interaction laws for small and large waves in the Ginzburg-Landau equation, *IMA Journal of Applied Mathematics* **46**, 121–136. Nonlinear Galerkin method and theory.

Pudykiewicz, J., Benoit, R. and Staniforth, A.: 1985, Preliminary results from a partial LRTAP model based on an existing meteorological forecast model, *Atmospheres and Oceans* **23**, 267–303. Not spectral; convergence of fixed point iteration for semi-Lagrangian methods.

Pulicani, J. P. and Ouazzani, J.: 1991, A Fourier-Chebyshev pseudospectral method for solving steady 3-D Navier-Stokes and heat equations in cylindrical cavities, *Computers and Fluids* **20**(2), 93–109.

Purnell, D. K.: 1976, Solution of the advective equation by upstream interpolation with a cubic spline, *Monthly Weather Review* **104**, 42–48.

Purser, R. J. and Leslie, L. M.: 1988, A semi-implicit semi-Lagrangian finite-difference scheme using high-order spatial differencing on a nonstaggered grid, *Monthly Weather Review* **116**, 2069–2080. Not spectral, but very high order spatial differences.

Purser, R. J. and Leslie, L. M.: 1991, An efficient interpolation procedure for high order three-dimensional semi-Lagrangian models, *Monthly Weather Review* **119**, 2492–2498. Not spectral, but high order spatial differences.

Purser, R. J. and Leslie, L. M.: 1994, An efficient semi-Lagrangian scheme using third-order semi-implicit time integration and forward trajectories, *Monthly Weather Review* **122**, 745–756.

Qin, M. Z. and Zhang, M. Q.: 1990, Multi-stage symplectic schems of two kinds of Hamiltonian systems for wave equations, *Computer Mathematics and Applications* **19**, 51–62.

Rand, R. H.: 1984, *Computer Algebra in Applied Mathematics: An Introduction to MACSYMA*, Pitman Research Notes in Mathematics, Pitman, New York.

Rand, R. H. and Armbruster, J.: 1987, *Perturbation Methods, Bifurcation Theory and Computer Algebra*, Springer-Verlag, New York. Not spectral, but many of the perturbation series are spectral series, and the order-by-order equations are solved by Galerkin methods.

Randriamampianina, A.: 1994, On the use of vorticity-vector-potential with a spectral tau method in rotating annular domains, *in* C. Bernardi and Y. Maday (eds), *Analysis, Algorithms and Applications of Spectral and High Order Methods for Partial Differential Equations*, Selected Papers from the International Conference on Spectral and High Order Methods (ICOSAHOM '92), Le Corum, Montpellier, France, 22-26 June 1992, North-Holland, Amsterdam, pp. 439–448. Three-dimensional vector potential for multiply connected domain with Fourier in aximuthal coordinate and Chebyshev in the radial and axial coordinates with time discretization by a mixed second order Adams-Bashforth/second order Backwards Euler [Backward Differentiation] . Also published in Finite Elements in Analysis and Design, vol. 16, pp. 299-307.

Randriamampianina, A., Bontoux, P. and Roux, B.: 1987, Ecoulements induits par la force gravique dans une cavité cylindrique en rotation, *International Journal of Heat and Mass*

Transfer **30**, 1275–1292. Spectral tau-method study of the competition between buoyancy and rotational forces in a confined cavity.

Rančić, M., Purser, R. J. and Mesinger, F.: 1996, A global shallow water model using an expanded spherical cube: Gnomonic versus conformal coordinates, *Quart. J. Roy. Meteor. Soc.* **122**, 959–982. Finite difference model: alternative to spherical harmonics method.

Rasch, P. and Williamson, D.: 1990a, On shape-preserving interpolation and semi-Lagrangian transport, *SIAM Journal of Scientific and Statistical Computing* **11**, 656–687.

Rasch, P. and Williamson, D.: 1990b, Computational aspects of moisture transport in global models of the atmosphere, *Quarterly Journal of the Royal Meteorological Society* **116**, 1071–1090.

Rasch, P. and Williamson, D.: 1991, The sensitivity of a general circulation model climate to the moisture transport formulation, *Journal of Geophysical Research* **96**, 13123–13137.

Rashid, A., Cao, W.-M. and Guo, B.-Y.: 1993, Three level Fourier pseudospectral method for fluid flow with low Mach number, *Journal of SUST* **16**, 223–237.

Rashid, A., Cao, W.-M. and Guo, B.-Y.: 1994a, Three level Fourier spectral approximations for fluid flow with low Mach number, *AMC* **63**, 131–150.

Rashid, A., Cao, W.-M. and Guo, B.-Y.: 1994b, The Fourier pseudospectral method for fluid flow with low Mach number, *Journal of Applied Science* **12**, 1–12.

Raspo, I., Ouazzani, J. and Peyret, R.: 1994, A direct Chebyshev multidomain method for flow computation with application to rotating systems, *Contemporary Mathematics* **180**, 533–538. 2D axisymmetric Navier-Stokes flow in a cylinder in vorticity/stream function form with influence matrix solution; two domains with a singularity in the corner of one domain wall where the boundary conditions change; fast algebraic rate of convergence.

Raspo, I., Ouazzani, J. and Peyret, R.: 1996, A spectral multidomain technique for the computation of the Czochralski melt configuration, *International Journal for Numerical Methods in Fluid Mechanics* **6**, 31–58. Similar to previous article, but different engineering application.

Reid, J. K. and Walsh, J. E.: 1965, An elliptic eigenvalue problem for a re-entrant region, *SIAM Journal of Applied Mathematics* **13**, 837–850. Devised a global mapping employed by Mason (1967); not spectral.

Rennich, S. C. and Lele, S. K.: 1997, Numerical method for incompressible vortical flows with two unbounded directions, *J. Comput. Phys.* **137**, 101–129. Three-dimensional Fourier method with domain truncation, using analytical conditions on the irrotational part of the velocity to greatly reduce the size of the domain.

Richardson, L. F.: 1910, The approximate arithmetical solution by finite differences of physical problems involving differential equations with an application to stresses in a masonry dam, *Phil. Trans. Royal Society of London* **A210**, 307–357. Invention of Richardson's iteration.

Richardson, L. F.: 1922, *Weather Prediction by Numerical Processes*, Cambridge University Press, New York. Ur-text of computational weather forecasting.

Riley, J. J., Metcalfe, R. W. and Orszag, S. A.: 1986, Direct numerical simulations of chemi-
    cally reacting turbulent mixing layers, *Physics of Fluids* **29**, 406–422.

Ritchie, H.: 1986, Eliminating the interpolation associated with the semi-Lagrangian
    scheme, *Monthly Weather Review* **114**(1), 135–146. Not spectral, but the time-marching
    algorithm was applied to spectral models in several of the author's later articles.

Ritchie, H.: 1987, Semi-Lagrangian advection on a Gaussian grid, *Monthly Weather Review*
    115, 608–619. Advection by a steady wind in spherical geometry.

Ritchie, H.: 1988, Application of the semi-Lagrangian method to a spectral model of the
    shallow-water equations, *Monthly Weather Review* **116**, 1587–1598. Surface of a sphere.

Ritchie, H.: 1991, Application of the semi-Lagrangian method to a multilevel spectral prim-
    itive equations model, *Quart. J. Roy. Met. Soc.* **117**, 91–106.

Ritchie, H. and Tanguay, M.: 1996, A comparison of spatially averaged Eulerian and semi-
    Lagrangian treatment of mountains, *Monthly Weather Review* **124**, 167.

Robert, A.: 1981, A stable numerical integration scheme for the primitive meteorological
    equations, *Atmosphere—Oceans* **19**, 35–46.

Robert, A.: 1982, A semi-Lagrangian and semi-implicit numerical integration scheme for
    the primitive meteorological equations, *Journal of the Meteorological Society of Japan*
    60, 319–324.

Robert, A. J.: 1966, The integration of a low order spectral form of the primitive meteoro-
    logical equations, *J. Meteor. Soc.* **44**, 237–244. Alternatives to spherical harmonics. First
    use of what is now called the "Asselin" time filter to damp high temporal frequencies.

Robert, A., Yee, T. L. and Ritchie, H.: 1985, A semi-Lagrangian and semi-implicit numer-
    ical integration scheme for multilevel atmospheric models, *Monthly Weather Review*
    113, 388–394.

Robinson, A. C. and Saffman, P. G.: 1984, Stability and structure of stretched vortices, *Stud-
    ies in Applied Mathematics* **68**, 163–181. Sinc series: 2D, nonlinear eigenvalue problem
    on an unbounded domain.

Robson, R. E. and Prytz, A.: 1993, The discrete ordinate/pseudo-spectral method: Review
    and application from a physicist's perspective, *Australian J. Physics* **46**, 465–495. RE-
    VIEW.

Rogallo, R. S.: 1977, An ILLIAC program for the numerical simulation of homogeneous,
    incompressible turbulence, *Technical Report TM-73203*, NASA.

Rogallo, R. S.: 1981, Numerical experiments in homogeneous turbulence, *Technical Report
    TM-81315*, NASA.

Rogallo, R. S. and Moin, P.: 1984, Numerical simulation of turbulent flows, *Annual Reviews
    of Fluid Mechanics* **16**, 99–137.

Ronchi, C., Iacono, R. and Paolucci, P. S.: 1996, The "Cubed Sphere": A new method for
    the solution of partial differential equations in spherical geometry, *J. Comput. Phys.*
    124(1), 93–114. Comparison of new finite difference method with spherical harmonics.

Rønquist, E. M.: 1988, *Optimal spectral element methods for the unsteady three-dimensional incompressible Navier-Stokes equations*, PhD dissertation, MIT, Department of Mechanical Engineering.

Rønquist, E. M.: 1996, Convection treatment using spectral elements of different order, *Int. J. Numer. Meth. Fluids* **22**(4), 241–264.

Rønquist, E. M. and Patera, A. T.: 1987a, A Legendre spectral element method for the Stefan problem, *Int. J. Num. Methods. Eng.* **24**, 2273–2299.

Rønquist, E. M. and Patera, A. T.: 1987b, Spectral element multigrid. I. Formulation and numerical results, *Journal of Scientific Computing* **2**, 389–406.

Runge, C.: 1903, *Zeit. f. Math. u. Phys.* **48**, 443–456.

Runge, C.: 1905, *Zeit. f. Math. u. Phys.* **53**, 117–123.

Runge, C.: 1913, *Berlauterung des Rechnungsformulars, u. s. w., Braunschweig.* These three papers describe Runge's independent discovery of what is now called the Fast Fourier Transform.

Saff, E. B. and Kuijlaars, A. B. J.: 1997, Distributing many points on a sphere, *Mathematical Intelligencer* **19**(1), 5–11. Semi-popular review of progress and open problems in generating nearly uniform distributions of points, such as collocation points, on a sphere.

Saffman, P. G. and Szeto, R.: 1981, Structure of a linear array of uniform vortices, *Studies in Applied Mathematics* **65**, 223–248. Application of sinc spectral method.

Salupere, A., Maugin, G. A., Engelbrecht, J. and Kalda, J.: 1996, On the KdV soliton formation and discrete spectral analysis, *Wave Motion* **23**, 49–66. High resolution Fourier solutions in one space dimension.

Sanders, B. F., Katopodes, N. and Boyd, J. P.: 1998, Unified pseudo-spectral solution to water wave equations, *Journal of Hydraulic Eng. ASCE* **124**, 2–12. Fourier pseudospectral calculations of a wide variety of one-dimensional nonlinear wave equations with careful comparisons with exact soliton solutions.

Sanugi, B. B. and Evans, D. J.: 1988, A Fourier series method for the numerical solution of the nonlinear advection problem, *Applied Mathematics Letters* **1**, 385–389.

Sanz-Serna, J. M. and Calvo, M. P.: 1994, *Numerical Hamiltonian Methods*, Chapman and Hall, London.

Sardeshmukh, P. D. and Hoskins, B. J.: 1984, Spatial smoothing on the sphere, *Monthly Weather Review* **112**, 2524–2529. If a filter applied to a spherical harmonics series depends only on the subscript n of the spherical harmonics, it is equivalent to an isotropic spatial filter whose effect depends only on distance from the point of observation.

Sawyer, J. S.: 1963, A semi-Lagrangian method of solving the vorticity equation, *Tellus* **15**(4), 336–342. Not spectral; coined the label "semi-Lagrangian".

Schaffer, S. and Stenger, F.: 1986, Multigrid-sinc methods, *Applied Mathematics and Computation* **19**, 311–319.

Schatz, M. F., Tagg, R. P., Swinney, H. L., Fischer, P. F. and Patera, A. T.: 1991, Supercritical transition in plane channel flow with spatially periodic perturbations, *Physical Review Letters*.

Schmidt, F.: 1977, Variable fine mesh in the spectral global models, *Beitr. Phys. Atmos.* **50**, 211–217. Conformal mapping to give high resolution in a chosen region.

Schmidt, F.: 1982, Cyclone tracing, *Beitr. Phys. Atmos.* **55**, 335–357. Conformal mapping is used to track cyclones by increasing resolution locally around the track of the storm; method of Schmidt(1977).

Schnack, D. D., Baxter, D. C. and Caramana, E. J.: 1984, A pseudospectral algorithm for three-dimensional magnetohydrodynamic simulation, *Journal of Computational Physics* **55**, 485–514.

Schneidesch, C., Deville, M. and Demaret, P.: 1990, Steady-state solution of a convection benchmark problem by multidomain Chebyshev collocation, *in* B. Roux (ed.), *Numerical Simulation of Oscillatory Convection in Low-Pr Fluids*, Friedr. Vieweg & Sohn Verlagsgesellschaft mbH, Braunschweig, part 4, pp. 256–261.

Schneidesch, C. R. and Deville, M. O.: 1993, Chebyshev collocation method and multidomain decomposition for Navier-Stokes equations in complex curved geometries, *Journal of Computational Physics* **106**, 234–257.

Schneidesch, C. R. and Deville, M. O.: 1994, Multidomain decomposition of curved geometries in the Chebyshev collocation method for thermal problems, *in* C. Bernardi and Y. Maday (eds), *Analysis, Algorithms and Applications of Spectral and High Order Methods for Partial Differential Equations*, Selected Papers from the International Conference on Spectral and High Order Methods (ICOSAHOM '92), Le Corum, Montpellier, France, 22-26 June 1992, North-Holland, Amsterdam, pp. 87–94. Also in Comput. Meths. Appl. Mech. Engrg., 116.

Schoeberl, M. R. and Geller, M. A.: 1977, A calculation of the structure of stationary planetary waves in winter, *Journal of the Atmospheric Sciences* **34**, 1235–1255. Application of finite differences and block-tridiagonal Gaussian elimination to a PDE of mixed elliptic-hyperbolic type.

Schultz, W. W., Huh, J. and Griffin, O. M.: 1994, Potential-energy in steep and breaking waves, *J. Fluid Mech.* **278**, 201–228. Spectral algorithm for two-dimensional potential flow.

Schultz, W. W., Lee, N.-Y. and Boyd, J. P.: 1989, Chebyshev pseudospectral method of viscous flows with corner singularities, *Journal of Scientific Computing* **4**, 1–24.

Schultz, W. W., Vanden-Broeck, J., Jiang, L. and Perlin, M.: 1998, Highly nonlinear standing water waves with small capillary effect, *J. Fluid Mech.* **369**, 253–272. Spectral boundary integral method is combined with Newton iteration to compute spatially and temporally periodic standing waves. A non-uniform node distribution (change-of-coordinate) is used to get accurate approximations near the limiting wave.

Schumack, M. R.: 1996, Application of the pseudospectral method to thermodynamic lubrication, *Int. J. Numer. Meth. Fluids* **23**(11), 1145–1161. Preprint.

Schumack, M. R., Schultz, W. W. and Boyd, J. P.: 1991, Spectral method solution of the Stokes equations on nonstaggered grids, *Journal of Computational Physics* **94**(1), 30–58.

Schumer, J. W. and Holloway, J. P.: 1998, Vlasov simulations using velocity-scaled Hermite representations, *J. Comput. Phys.* **144**(2), 626–661. Hermite function calculations in plasma physics.

Schuster, A.: 1903, On some definite integrals and a new method of reducing a function of spherical coordinates to a series of spherical harmonics, *Phil. Trans. Roy. Soc. London A* **200**, 181–223. Two-step transform in which data for a given longitudinal wavenumber is expanded first as a trigonometric series, and then this is converted to spherical harmonics at a savings of roughly a factor of two over direct computation of spherical harmonic coefficients.

Schwartz, C.: 1963, Estimating convergence rates of variational calculations, *Methods in Computational Physics* **2**, 241–266. How singularities affect the rate of convergence of Laguerre function series.

Sela, J.: 1980, Spectral modelling at the National Meteorological Center, *Monthly Weather Review* **108**, 1279–1292.

Sela, J. G.: 1995, Weather forecasting on parallel architectures, *Parallel Comput.* **21**(10), 1639–1654. Spherical harmonic modelling at the National Meteorological Center.

Shen, J.: 1994, Efficient spectral-Galerkin method I. Direct solvers of second- and fourth-order equations using Legendre polynomials, *SIAM Journal of Scientific Computing* **15**(6), 1489–1505.

Shen, J.: 1995a, Efficient spectral-Galerkin method II. Direct solvers of second- and fourth-order equations using Chebyshev polynomials, *SIAM Journal of Scientific Computing* **16**(6), 74–87.

Shen, J.: 1995b, On fast Poisson solver, inf-sup constant and iterative Stokes solver by Legendre Galerkin method, *Journal of Computational Physics* **116**, 184–188.

Shen, J.: 1997, Efficient spectral-Galerkin method III. Polar and cylindrical geometries, *SIAM J. Sci. Comput.* Fast methods for inverting the Laplace and other almost constant coefficient operators using orthogonal polynomials.

Sherwin, S. J.: 1997, Hierarchical hp finite elements in hybrid domains, *Finite Elements in Analysis and Design* **27**, 109–119.

Sherwin, S. J. and Karniadakis, G. E.: 1996, Tetrahedral hp finite elements: algorithms and flow simulations, *Journal of Computational Physics* **124**, 14–45. Tensor product spectral elements in three dimensions.

Sherwin, S. J., Karniadakis, G. E. and Orszag, S. A.: 1994, Numerical simulation of the ion etching process, *Journal of Computational Physics* **110**, 373.

Shoucri, M. M. and Gagné, R. R. J.: 1977, Numerical solution of a two-dimensional Vlasov equation, *Journal of Computational Physics* **25**, 94–103. Tensor product Hermite basis for the two velocity coordinates, finite differences for the two space dimensions.

Simmons, A. J. and Hoskins, B. J.: 1975, A comparison of spectral and finite-difference simulations of a growing baroclinic wave, *Quarterly Journal of the Royal Meteorological Society* **101**, 551–565.

Siyyam, H. I. and Syam, M. I.: 1997, An accurate solution of the Poisson equation by the Chebyshev-Tau method, *J. Comput. App. Math.* **85**, 1–10. Alternative to Haidvogel-Zang algorithm for separable PDEs.

Sloan, D. M.: 1991, Fourier pseudospectral solution of the Regularised Long Wave equation, *Journal of Computational and Applied Mathematics* **36**, 159–179.

Sloan, I. H.: 1995, Polynomial interpolation and hyperinterpolation over general regions, *Journal of Approximation Theory* **83**, 238–254. Computation of approximations on the surface of sphere as a truncated spherical harmonic series with coefficients evaluated by numerical quadrature. "Hyperinterpolation" is a coefficient computed by a quadrature with more points than unknowns in the truncated basis. For the sphere, interpolation is not possible but hyperinterpolation is successful.

Smith, N. R.: 1988, A truncated oceanic spectral model for equatorial thermodynamic studies, *Dyn. Atmos. Oceans* **12**, 313–337. Hermite spectral series.

Smith, R. C. and Bowers, K. L.: 1991, A fully Galerkin method for the recovery of stiffness and damping parameters in Euler-Bernoulli beam models, *Computation and Control II, Proceedings of the Bozeman Conference 1990*, number 11 in *Progress in Systems and Control Theory*, Birkhäuser, Boston, pp. 289–306.

Smith, R. C., Bogar, G. A., Bowers, K. L. and Lund, J.: 1991a, The sinc-Galerkin method for fourth-order differential equations, *SIAM Journal of Numerical Analysis* **28**, 760–788.

Smith, R. C., Bowers, K. L. and Lund, J.: 1991b, Efficient numerical solution of fourth-order problems in the modeling of flexible structures, *Computation and Control, Proceedings of the Bozeman Conference 1988*, number 1 in *Progress in Systems and Control Theory*, Birkhäuser, Boston, pp. 289–306.

Smith, R. C., Bowers, K. L. and Lund, J.: 1992, A fully sinc-Galerkin method for Euler-Bernoulli beam models, *Numerical Methods for Partial Differential Equations* **8**, 171–202.

Smolarkiewicz, P. and Rasch, P.: 1990, Monotone advection on the sphere: An Eulerian versus a semi-Lagrangian approach, *Journal of the Atmospheric Sciences* **48**, 793–810.

Sneddon, G. E.: 1996, Second-order spectral differentiation matrices, *SIAM Journal of Numerical Analysis* **33**(6), 2468–2487. Shows that different discretization methods (pseudospectral versus Galerkin versus tau) are equivalent, after splitting the matrix into two subproblems through parity, to a rank-one update to the symmetric differentiation matrix and a different rank-one update to the differentiation matrix for functions antisymmetric with respect to x. Claims that one can choose an update which makes the eigenvalues $O(N^2)$ at largest instead of $O(N^4)$ without drastic loss of accuracy in differentiation.

Solomonoff, A.: 1992, A fast algorithm for spectral differentiation, *Journal of Computational Physics* **98**, 174–177. Independent invention of the Parity Matrix Multiplication.

Spalart, P. R.: 1984, A spectral method for external viscous flows, *Contemporary Mathematics* **28**, 315–335. Divergence-free basis functions.

Spotz, W. F., Taylor, M. A. and Swarztrauber, P. N.: 1998, Fast shallow-water equation solvers in latitude-longitude coordinates, *Journal of Computational Physics* **145**(1), 432–444. Double Fourier series in both longitude and latitude, which would normally have serious pole troubles, are stabilized by using a filter. One filter is an improvement of a Fourier filter developed by Merilees. Another is the fast multipole-like filter develped by Jakob-Chien and Alpert and improved by Yarvin and Rokhlin. Both filters work well and eliminate most of the speed penalty associated with Legendre transforms in a pure spherical harmonics method.

Staniforth, A. and Côté, J.: 1991, Semi-Lagrangian integration schemes for atmospheric models — a review, *Monthly Weather Review* **119**, 2206–2223.

Stenger, F.: 1979, A "Sinc-Galerkin" method of solution of boundary value problems, *Mathematics of Computation* **33**, 85–109.

Stenger, F.: 1981, Numerical methods based on Whittaker cardinal or Sinc functions, *SIAM Review* **23**, 165–224.

Stenger, F.: 1993, *Sinc Methods*, Springer-Verlag, New York. 500 pp. Restricted to sinc functions; nothing on Chebyshev or Fourier algorithms.

Strain, J.: 1994, Fast spectrally-accurate solution of variable-coefficient elliptic problems, *Proceedings of the American Mathematical Society* **122**, 843–850. Iteration for non-separable BVPs in which a separable elliptic problem is solved by a fast direct spectral method at each iteration.

Strain, J.: 1995, Spectral methods for nonlinear parabolic systems, *Journal of Computational Physics* **122**, 1–12.

Straka, J. M., Wilhelmson, R. B., Wicker, L. J., Anderson, J. R. and Droegemeier, K. K.: 1993, Numerical solutions of a non-linear density current: A benchmark solution and comparisons, *International Journal for Numerical Methods in Fluids* **17**, 1–22.

Strang, G.: 1968, On the construction and comparison of difference schemes, *SIAM Journal of Numerical Analysis* **5**(3), 506–517. Independent invention of fractional steps time-marching, which is often called "Strang splitting" in the U. S.

Strang, G. and Fix, G. J.: 1973, *The Analysis of the Finite Element Method*, Prentice-Hall, Englewood Cliffs, New Jersey. 305 pp.

Straughan, B. and Walker, D. W.: 1996, Two very accurate and efficient methods for computing eigenvalues and eigenfunctions in porous convection problems, *Journal of Computational Physics* **127**, 128–141. Compound matrix and Chebyshev-tau algorithms.

Streett, C. L. and Zang, T. A.: 1984, Spectral methods for the solution of the boundary-layer equations, *Paper 84-0170*, AIAA. Nonlinear Richardson's iteration.

Streett, C. L., Zang, T. A. and Hussaini, M. Y.: 1985, Spectral multigrid methods with applications to transonic potential flow, *Journal of Computational Physics* **57**, 43–76.

Stuhne, G. R. and Peltier, W. R.: 1999, New icosahedral grid-point discretizations of the shallow water equations on the sphere, *J. Comput. Phys.* **148**(1), 23–58. Non-spectral alternative to spherical harmonics.

Su, Y. Y. and Khomami, B.: 1992, Numerical solution of eigenvalue problems using spectral techniques, *Journal of Computational Physics* **100**, 297–305.

Sulem, P. L., Sulem, C. and Patera, A.: 1985, A numerical simulation of singular solutions to the two-dimensional Cubic Schrodinger equation, *Communications in Pure and Applied Mathematics* **37**, 755–778.

Süli, E. and Ware, A.: 1991, A spectral method of characteristics for hyperbolic problems, *SIAM Journal of Numerical Analysis* **28**, 423–445.

Süli, E. and Ware, A. F.: 1992, Analysis of the spectral Lagrange-Galerkin method for the Navier-Stokes equations, *in* J. G. Heywood, K. Masuda, R. Rautmann and V. A. Solonnikov (eds), *The Navier-Stokes Equations II-Theory and Numerical Methods: Proceedings of a Conference in Oberwolfach, Germany*, Springer-Verlag, pp. 184–195.

Swarztrauber, P. N.: 1977, On the spectral approximation of discrete scalar and vector functions on the sphere, *SIAM Journal of Numerical Analysis* **16**, 934–949.

Swarztrauber, P. N.: 1981, The approximation of vector functions and their derivatives on the sphere, *SIAM Journal of Numerical Analysis* **18**, 191–210.

Swarztrauber, P. N.: 1982, Vectorizing the FFTs, *in* G. Rodrique (ed.), *Parallel Computations*, Academic Press, New York, pp. 51–83.

Swarztrauber, P. N.: 1986, Symmetric FFTs, *Mathematics of Computation* **47**, 323–346. Efficient methods of computing cosine and sine transforms.

Swarztrauber, P. N.: 1987, Multiprocessor FFTs, *Parallel Computing* **7**, 197–210.

Swarztrauber, P. N.: 1993, The vector harmonic transform method for solving partial differential equations in spherical geometry, *Monthly Weather Review* **121**, 3415–3437.

Swarztrauber, P. N.: 1996, Spectral transform methods for solving the shallow-water equations on the sphere, *Monthly Weather Review* **124**, 730.

Swarztrauber, P. N. and Kasahara, A.: 1985, The vector harmonic analysis of Laplace's tidal equations, *SIAM Journal of Scientific and Statistical Computing* **6**, 464–491. This and earlier individual papers by the two authors are very good works on the use of vector basis functions for geophysical flows.

Swarztrauber, P. N., Williamson, D. L. and Drake, J. B.: 1997, The Cartesian method for solving partial differential equations in spherical geometry, *Dyn. Atmos. Oceans* **27**, 679–706. Ingenious non-spectral scheme on an icosahedral grid.

Tan, B. and Boyd, J. P.: 1997, Dynamics of the Flierl-Petviashvili monopoles in a barotropic model with topographic forcing, *Wave Motion* **26**, 239–252. Double Fourier series for nonlinear initial value problem.

Tan, B. and Boyd, J. P.: 1998, Davydov soliton collisions, *Physics Letters A* **240**, 282–286. Double Fourier series for nonlinear initial value problem.

Tan, B. and Boyd, J. P.: 1999, Coupled-mode envelope solitary waves in a pair of cubic Schroedinger equations with cross modulation, *Nonlinearity*. Submitted.

Tan, C. S.: 1985, Accurate solution of three-dimensional Poisson's equation in cylindrical coordinates by expansion in Chebyshev polynomials, *Journal of Computational Physics* **59**, 81–95.

Tang, T.: 1993, The Hermite spectral method for Gaussian-type functions, *SIAM Journal of Scientific Computing* **14**, 594–606.

Taylor, M.: 1995, Cubature for the sphere and the discrete spherical harmonic transform, *SIAM Journal of Numerical Analysis* **32**, 667–670.

Taylor, M. A. and Wingate, B. A.: 1999, The Fekete collocation points for triangular spectral elements, *SIAM Journal of Numerical Analysis*. Submitted.

Taylor, M., Tribbia, J. and Iskandarani, M.: 1997, The spectral element method for the shallow water equations on the sphere, *Journal of Computational Physics* **130**(1), 92–108.

Taylor, T. D., Hirsh, R. S. and Nadworny, M. M.: 1981, FFT versus conjugate gradient method for solutions of flow equations by pseudospectral methods, *in* H. Viviand (ed.), *Proceedings of the 4th GAMM Conference on Numerical Methods in Fluid Mechanics*, Vieweg, Braunschweig, pp. 311–325.

Taylor, T. D., Hirsh, R. S. and Nadworny, M. M.: 1984, Comparison of FFT, direct inversion, and conjugate gradient methods for use in pseudospectral methods, *Computers and Fluids* **12**, 1–9. Examines the penalty for using a nonstandard grid such that neither the FFT nor matrix multiplication transform are applicable, and one must invert matrices to calculate coefficients of f(x) from grid point values of f(x).

Témam, R.: 1989, Attractors for the Navier-Stokes equations: localization and approximation, *Journal of the Faculty of Science of the University of Tokyo, Sec. Ia, Mathematics* **36**, 629–647.

Témam, R.: 1991a, New emerging methods in numerical analysis; applications to fluid mechanics, *in* M. Gunzberger and N. Nicolaides (eds), *Computational Fluid Dynamics — Trend and Advances*, Cambridge University Press, Cambridge.

Témam, R.: 1991b, Stability analysis of the nonlinear Galerkin method, *Mathematics of Computation* **57**, 477–503.

Témam, R.: 1992, General methods for approximating inertial manifolds. applications to computing, *in* D. S. Broomhead and A. Iserles (eds), *The Dynamics of Numerics and the Numerics of Dynamics*, Clarendon Press, Oxford, Oxford, pp. 1–21.

Temperton, C.: 1975, Algorithms for the solutions of cyclic tridiagonal systems, *Journal of Computational Physics* **19**, 317–323.

Temperton, C.: 1983a, Fast mixed-radix real Fourier transforms, *Journal of Computational Physics* **52**, 340–350.

Temperton, C.: 1983b, Self-sorting mixed-radix fast Fourier transforms, *Journal of Computational Physics* **52**, 1–23.

Temperton, C.: 1985, Implementation of a self-sorting in-place prime factor FFT algorithm, *Journal of Computational Physics* **58**, 283–299.

Temperton, C.: 1992, A generalized prime factor FFT algorithm for any N=(2**p) (3**q) (5**r), *SIAM J. Sci. Stat. Comput.* **13**, 676–686.

Temperton, C. and Staniforth, A.: 1987, An efficient two-time-level semi-Lagrangian semi-implicit integration scheme, *Quarterly Journal of the Royal Meteorological Society* **113**, 1025–1039.

Thompson, J., Warsi, Z. and Mastin, C.: 1985, *Numerical Grid Generation*, North-Holland, New York. Not spectral.

Thompson, W. D.: 1917, *Growth and Form*, Cambridge University Press, Cambridge. Not spectral, 600 pp.

Thuburn, J.: 1997, A PV-based shallow-water model on a hexagonal-icosahedral grid, *Mon. Wea. Rev.* **125**, 2328–2347. Non-spectral alternative for flows on the surface of a sphere.

Trefethen, L. N. and Bau, III, D.: 1997, *Numerical Linear Algebra*, Society for Industrial and Applied Mathematics (SIAM), Philadelphia.

Tribbia, J. J.: 1984a, Modons in spherical geometry, *Geophys. Astrophys. Fluid Dyn.* **30**, 131–168. Spherical harmonics in a rotated coordinate system are used to construct nonlinear Rossby waves.

Tribbia, J. J.: 1984b, A simple scheme for high-order nonlinear normal initialization, *Monthly Weather Review* **112**, 278–284. Iterative scheme for initialization onto the slow manifold, illustrated by Hough-Hermite spectral basis.

Tse, K. L. and Chasnov, J. R.: 1998, A Fourier-Hermite pseudospectral method for penetrative convection, *Journal of Computational Physics* **142**(2), 489–505. Two Fourier coordinates plus Hermite functions in the unbounded vertical; motion is confined to an unstable layer far from any boundaries.

Tuckerman, L.: 1989, Divergence-free velocity field in nonperiodic geometries, *Journal of Computational Physics* **80**, 403–441. Very careful treatment of influence matrix method and spurious pressure modes in computing incompressible flow.

Vallis, G. K.: 1985, On the spectral integration of the quasi-geostrophic equations for doubly-periodic and channel flow, *Journal of the Atmospheric Sciences* **42**, 95–99.

Van Kemenade, V. and Deville, M.: 1994a, Preconditioned Chebyshev collocation for non-Newtonian fluid flows, *in* C. Bernardi and Y. Maday (eds), *Analysis, Algorithms and Applications of Spectral and High Order Methods for Partial Differential Equations*, Selected Papers from the International Conference on Spectral and High Order Methods (ICOSAHOM '92), Le Corum, Montpellier, France, 22-26 June 1992, North-Holland, Amsterdam, pp. 377–386. Also in Finite Elements in Analysis and Design, vo. 16, pp. 237-245.

Van Kemenade, V. and Deville, M. O.: 1994b, Application of spectral elements to viscoelastic creeping flows, *Journal of Non-Newtonian Fluid Mechanics* **51**, 277–308. Plane and cylindrical geometries; Maxwell-B fluid; comparisons with 4 x 4 SUPG finite elements.

van Loan, C.: 1992, *Computational Frameworks for the Fast Fourier Transform*, SIAM, Philadelphia.

Vandeven, H.: 1991, Family of spectral filters for discontinuous problems, *SIAM Journal of Scientific Computing* **6**, 159–192.

variational spectral method for the two-dimensional Stokes problem, A.: 1998, S.-r. jun and y.-h. kwon and s. kang, *Computers Math. Applic.* **35**(4), 1–17.

Verkley, W. T. M.: 1997a, A pseudo-spectral model for two-dimensional incompressible flow in a circular basin. I. Mathematical formulation, *Journal of Computational Physics* **136**(1), 100–114. Employs the basis set $r^{|m|} P_k^{0,|m|}(2r^2 - 1) \exp(im\theta)$ where $P_k^{\alpha,\beta}$ is the usual Jacobi polynomial; shows and exploits the fact that the inverse of Laplace operator has a banded Galerkin representation with this basis.

Verkley, W. T. M.: 1997b, A pseudo-spectral model for two-dimensional incompressible flow in a circular basin. II. Numerical examples, *Journal of Computational Physics* **136**(1), 115–131.

Vermer, J. G. and van Loon, M.: 1994, An evaluation of explicit pseudo-steady-state approximation schemes to stiff ODE systems from chemical kinetics, *Journal of Computational Physics* **113**, 347–352. The PSSA strategy, widely used in chemical kinetics, is equivalent to the lowest order Nonlinear Galerkin method: the ODEs for the fast components are replaced by algebraic relations.

Vianna, M. L. and Holvorcem, P. R.: 1992, Integral-equation approach to tropical ocean dynamics. 1. Theory and computational methods, *J. Marine Res.* **50**(1), 1–31. Boundary-element algorithm constructed through ingenious summation of slowly converging Hermite series.

Vinograde, B.: 1967, *Linear and Matrix Algebra*, D. C. Heath, Boston. Not spectral.

Voigt, R. G., Gottlieb, D. and Hussaini, M. Y. (eds): 1995, *Spectral Methods for Partial Differential Equations*, SIAM, Philadelphia.

Vozovoi, L., Israeli, M. and Averbuch, A.: 1996, Analysis and application of the Fourier-Gegenbauer method to stiff differential equations, *SIAM J. Numer. Anal.* **33**(5), 1844–1863.

Vozovoi, L., Weill, A. and Israeli, M.: 1997, Spectrally accurate solution of non-periodic differential equations by the Fourier-Gegenbauer method, *SIAM J. Numer. Anal.* **34**(4), 1451–1471.

Wahba, G.: 1990, *Spline Models for Observational Data*, Vol. 59 of *CBMS-NSF Regional Conference Series in Applied Mathematics*, SIAM, Philadelphia. 169 pp. Fourier and spherical harmonic splines.

Wang, D.: 1991, Semi-discrete Fourier spectral approximations of infinite dimensional Hamiltonian systems and conservation laws, *Computers Mathematics and Applications* **21**, 63–75.

Ware, A. F.: 1991, *A spectral Lagrange-Galerkin method for convection-dominated diffusion problems*, Ph.D. dissertation, Oxford University, Wolfson College. 106 pp.

Ware, A. F.: 1994, A spectral Lagrange-Galerkin method for convection-dominated diffusion problems, *in* C. Bernardi and Y. Maday (eds), *Analysis, Algorithms and Applications of Spectral and High Order Methods for Partial Differential Equations*, Selected Papers from the International Conference on Spectral and High Order Methods (ICOSAHOM '92), Le Corum, Montpellier, France, 22-26 June 1992, North-Holland, Amsterdam, pp. 227–234. Also in Comp. Meths. Appl. Mech. Engrg., 116.

Ware, A. F.: 1998, Fast approximate Fourier transforms for irregularly spaced data, *SIAM Rev.* **40**(4), 838–856.

Weideman, J. A. C.: 1992, The eigenvalues of Hermite and rational spectral differentiation matrices, *Numerische Mathematik* **61**, 409–431.

Weideman, J. A. C.: 1994a, Computation of the complex error function, *SIAM Journal of Numerical Analysis* **31**, 1497–1518. Errata: 1995, 32, 330–331]. These series of rational functions are useful for complex-valued z.

Weideman, J. A. C.: 1994b, Computing integrals of the complex error function, *Proceedings of Symposia in Applied Mathematics* **48**, 403–407. Short version of Weideman(1994a).

Weideman, J. A. C.: 1995a, Errata: computation of the complex error function, *SIAM Journal of Numerical Analysis* **32**, 330–331.

Weideman, J. A. C.: 1995b, Computing the Hilbert Transform on the real line, *Mathematics of Computation* **64**(210), 745–762. The Hilbert transform of a function f is defined by $H\{f\}(y) \equiv \frac{1}{\pi}PV \int_{-\infty}^{\infty} \frac{f(x)}{x-y}dx$ where PV is the principal value of the (singular) integral. Weideman shows that a set of complex-valued orthogonal functions, $\rho_n(x) = (l + ix)^n/(l - ix)^{n+l}, n = 0, \pm 1, \pm 2, \ldots$, are eigenfunctions of the transform. Weideman's algorithm using these functions is generally more efficient than the Fourier transform algorithm, $H\{f\}(y) = F^{-1}\{i\text{sign}(k)F\{f\}(k)\}$. The Hilbert transform is important in many physical applications, notably the Benjamin-Ono model for solitary waves in deep stratified fluid.

Weideman, J. A. C. and Cloot, A.: 1990, Spectral methods and mappings for evolution equations on the infinite line, *Computer Methods in Applied Mechanics and Engineering* **80**, 467–481.

Weideman, J. A. C. and James, R. L.: 1992, Pseudospectral methods for the Benjamin-Ono equation, *Advances in Computer Methods for Partial Differential Equations VII* pp. 371–377. Both Fourier series and rational functions.

Welander, P.: 1955, Studies on the general development of motion in a two dimensional fluid, *Tellus* **7**, 141–156. Not spectral; classic article that shows (i) a square blob of fluid becomes greatly stretched and deformed even by very smooth flow — becoming "chaotically" mixed and bounded by a "fractal" curve, though these terms were not yet invented which (ii) shows the unfeasibility of a fully Lagrangian (as opposed to semi-Lagrangian) time-marching for fluids.

Weyl, H.: 1952, *Symmetry*, Princeton University Press, Princeton, New Jersey. Popular but rigorous discussion of symmetry in art and nature and its mathematical embodiment in group theory.

White, Jr., A. B.: 1982, On the numerical solution of initial/boundary value problems in one space dimension, *SIAM Journal of Numerical Analysis* **19**(4), 683–697. Shocks in Burgers' equation are resolved by transforming the spatial coordinate to arclength, which requires simultaneously integrating an extra equation.

Whittaker, E. T.: 1915, On the functions which are represented by the expansions of the interpolation theory, *Proceedings of the Royal Society Edinburgh* **35**, 181–194. Pioneering paper on sinc functions.

Wiin-Nielsen, A.: 1959, On the application of trajectory methods in numerical forecasting, *Tellus* **11**(2), 180–196. Not spectral; earliest application of semi-Lagrangian time-marching in which the choice of the particles is reinitialized at each time step to those arriving at the points of the regular Eulerian-coordinates grid at that time level.

Williamson, D.: 1990, Semi-Lagrangian transport in the NMC spectral model, *Tellus A* **42**, 413–428.

Williamson, D. L. and Rasch, P. J.: 1989, Two-dimensional semi-Lagrangian transport with shape-preserving interpolation, *Monthly Weather Review* **117**, 102–129. Both Cartesian and Gaussian (spherical) grid; only pure advection is computed.

Wingate, B. A. and Boyd, J. P.: 1996, Spectral element methods on triangles for geophysical fluid dynamics problems, *in* A. V. Ilin and L. R. Scott (eds), *Proceedings of the Third International Conference on Spectral and High Order Methods*, Houston Journal of Mathematics, Houston, Texas, pp. 305–314.

Wingate, B. A. and Taylor, M. A.: 1999a, A Fekete point triangular spectral element method; application to the shallow water equations, *Journal of Computational Physics*. Submitted.

Wingate, B. A. and Taylor, M. A.: 1999b, The natural function space for triangular spectral elements, *SIAM Journal of Numerical Analysis*. Submitted.

Wragg, A.: 1966, The use of Lanczos-τ methods in the numerical solution of a Stefan problem, *Computer Journal* **9**, 106–109.

Wunsch, C., Haidvogel, D. B., Iskandarani, M. and Hughes, R.: 1997, Dynamics of the long-period tides, *Progress in Oceanography* **40**(1–4), 81–108. Spectral elements.

Xu, C. and Maday, Y.: 1998, A spectral element method for the time-dependent two-dimensional Euler equations: applications to flow simulations, *J. Comput. Appl. Math.* **91**, 63–85. Show that a spectral approximation with polynomials of equal degree for pressure and velocity is well-posed. (Not true for the Navier-Stokes equation.).

Yakhot, A., Orszag, S. A., Yakhot, V. and Israeli, M.: 1989, Renormalization group formulation of large-eddy simulations, *Journal of Scientific Computing* **4**(2), 139–158.

Yanenko, N. N.: 1971, *The Method of Fractional Steps*, Springer-Verlag, New York. 160 pages.

Yang, G., Belleudy, P. and Temperville, A.: 1991, A higher-order Eulerian scheme for coupled advection-diffusion transport, *International Journal for Numerical Methods in Fluids* **12**, 43–58.

Yang, J. and Hsu, E.: 1991, On the use of the reach-back characteristics method for the calculation of dispersion, *International Journal for Numerical Methods in Fluids* **12**, 225–235.

Yang, T.-S. and Akylas, T. R.: 1996, Weakly nonlocal gravity-capillary solitary waves, *Physics of Fluids* **8**(6), 1506–1514. Rational Chebyshev functions (in somewhat unorthodox notation) and a special radiation basis function are combined with Newton's iteration to compute oscillatory tails even when the tails are smaller than 10^{-13} by combining collocation with Newton's iteration.

Yang, Z.-H. and Keller, H. B.: 1986, Multiple laminar flows through curved pipes, *Applied Numerical Mathematics* **2**, 257–271. The Dean problem of viscous steady flow through a coiled circular pipe is studied using Fourier expansion in polar angle and finite differences; good discussion of pseudoarclength continuation to trace an intricate structure of solution branches.

Yarvin, N. and Rohklin, V.: 1998, A generalized one-dimensional fast multipole method with application to filtering of spherical harmonics, *J. Comput. Phys.* **147**(2), 594–609.

Yee, S. Y. K.: 1981, Solution of Poisson's equation on a sphere by double Fourier series, *Monthly Weather Review* **109**, 501–505.

Yin, G.: 1994, Sinc-collocation method with orthogonalization for singular Poisson-like problems, *Mathematics of Computation* **62**, 21–40.

Young, D. M. and Gregory, R. T.: 1972, *A Survey of Numerical Mathematics, Vol. 1*, Addison-Wesley, Reading, MA. pg. 329–339; not spectral.

Z.-S. She, Jackson, E. and Orszag, S. A.: 1990, Vortex structure and dynamics in turbulence, *in* C. Canuto and A. Quarteroni (eds), *Spectral and High Order Methods for Partial Differential Equations: Proceedings of the ICOSAHOM '89 Conference in Como, Italy*, North-Holland/Elsevier, Amsterdam, pp. 173–184. Also in Comput. Meths. Appl. Mech. Engrg., vol. 80, with the same page numbers. Spectral computations of turbulence.

Zampieri, E. and Tagliani, A.: 1997, Numerical approximation of elastic waves by implicit spectral methods, *Comput. Meth. Appl. M.* **144**(1–2), 33–50. Legendre polynomials in single-domain spectral element (variational) algorithm combined with implicit time-marching; absorbing boundaries are used to imitate an infinite domain.

Zang, T.: 1991, On the rotation and skew-symmetric forms for incompressible flow simulations, *Applied Numerical Mathematics* **7**, 27–40.

Zang, T. A. and Hussaini, M. Y.: 1986, On spectral multigrid methods for the time-dependent Navier-Stokes equations, *Applied Mathematics and Computation* **19**, 359–372.

Zang, T. A., Krist, S. E. and Hussaini, M. Y.: 1989, Resolution requirements for numerical simulations of transition, *Journal of Scientific Computing* **4**(2), 197–219.

Zang, T. A., Wong, Y. S. and Hussaini, M. Y.: 1984, Spectral multigrid methods for elliptic equations II, *Journal of Computational Physics* **54**, 489–507.

Zebib, A.: 1984, A Chebyshev method for the solution of boundary value problems, *Journal of Computational Physics* **53**, 443–455. Assumes expansion for highest derivative of unknown rather than u itself.

Zebib, A.: 1987a, Stability of viscous flow past a circular cylinder, *Journal of Engineering Mathematics* **21**, 155–165.

Zebib, A.: 1987b, Removal of spurious modes encountered in solving stability problems by spectral methods, *Journal of Computational Physics* **70**(2), 521–525.

Zhao, S. and Yedlin, M. J.: 1994, A new iterative Chebyshev spectral method for solving the elliptic equation $\nabla \otimes (\sigma \nabla u) = f$, *Journal of Computational Physics* **113**, 215–223. The Haidvogel-Zang fast direct method for solving separable PDEs through Chebyshev series is the key operation in a fast iterative method for non-separable elliptic equations.

Zheng, J.-D., Zhang, R.-F. and Guo, B.-Y.: 1989, The Fourier pseudo-spectral method for the SRLW equation, *Applied Mathematics and Mechanics* **10**, 843–852.

Zrahia, U., Orszag, S. A. and Israeli, M.: 1996, Mixed analytical/numerical-spectral element algorithm for efficient solution of problems with interior or boundary layers, *in* A. V. Ilin and L. R. Scott (eds), *Proceedings of the Third International Conference on Spectral and High Order Methods*, Houston Journal of Mathematics, Houston, Texas, pp. 315–326. Analytical solution for boundary layer combined with spectral element solution for the outer flow.

1991 1985 1996 1986 1980 1965 1991 1989 1991 1996 1973 1995 1966 1967 1983
1989 1990 1992 1998a 1995 1998b 1997 1979 1981 1977 1977 1978 1975 1992 1978
1978 1986 1958 1986 1984 1982 1990 1985 1987 1992 1992 1994 1995 1989 1990 1962
1997 1978 1990 1992 1992 1996 1989 1991 1970 1984 1998 1996 1996 1970 1984 1994
1997 1996 1991 1997 1972 1974 1977 1995 1987 1978a 1978b 1978c 1980a 1980b 1980c
1981a 1981b 1981c 1982a 1982b 1982c 1982d 1982e 1983a 1983b 1983c 1983d 1984a
1984b 1984c 1984d 1984e 1984f 1984g 1985a 1985b 1985c 1985d 1986a 1986b 1987a
1987b 1987c 1987d 1988a 1988b 1988c 1989a 1989b 1989c 1989d 1989e 1990a 1990b
1990c 1990d 1991a 1991b 1991c 1991d 1991e 1991f 1992a 1992b 1992c 1992d 1992e
1993 1994a 1994b 1994c 1994d 1994e 1994f 1995a 1995b 1995c 1995d 1995e 1995f
1996a 1996b 1996c 1996d 1997a 1997b 1997c 1997d 1998a 1998b 1998c 1998d 1998e
1999a 1999b 1999c 2001 1982 1983 1987 1999 1991 1990 1998 1998 1999 1983 1985
1998 1986 1980 1987 1995 1981 1995 1994 1988 1992a 1989 1992b 1984 1967 1988 1985
1988 1994 1996 1996 1990 1968 1914 1985 1984 1990 1990 1970 1993a 1993 1993b
1982 1990 1988 1997 1957 1960 1997 1991 1990 1992 1990 1973 1965 1995 1990 1995
1961 1988 1994 1991 1996 1994 1985 1998 1987 1980 1981 1991 1978 1975 1984 1998
1994 1989 1991 1995 1980 1976 1977a 1977b 1977c 1979 1981a 1979 1971 1980 1981b
1986 1981 1989 1991 1989 1995 1995 1990 1982 1984 1985 1991 1981 1984 1990 1992
1994 1992 1933 1997 1998 1990 1994 1995 1997 1996 1974 1982 1995 1997 1998 1990
1998 1991 1993 1995 1961 1972 1987 1989 1992 1991 1983 1993 1994a 1994b 1997
1993 1974 1984 1966 1963 1987 1984 1989 1992 1973 1990 1994a 1994b 1997 1998
1997 1989 1991 1992 1994 1994 1988a 1989 1988b 1952 1955 1992 1987 1998 1991
1994 1988 1976 1975 1977 1978 1987 1988a 1988b 1990a 1990b 1992 1995 1996 1998
1997 1994 1978 1997 1973 1962 1968 1973 1972 1990 1966 1997 1992 1990 1986 1987a
1987b 1988a 1988b 1990a 1990b 1990c 1991 1992a 1992b 1993a 1993b 1994a 1994b
1997a 1997b 1988 1991 1990 1991 1989 1993 1988 1990 1984 1998 1995 1996 1995
1989 1970 1996 1997 1998 1972a 1972b 1972c 1986a 1986b 1993 1995 1996 1998 1993
1984 1988 1997 1985 1995 1996a 1996b 1997a 1997b 1995 1993 1995 1987 1982 1981
1984 1989 1983a 1983b 1977 1980 1994 1995a 1995b 1990 1985 1991 1993 1980 1985
1984a 1981a 1984b 1987a 1987b 1981b 1992 1967 1994 1993 1991 1993 1977 1990 1988
1998 1985 1996 1996 1977 1983 1979 1980 1981 1984 1980 1999 1997 1968 1994 1993
1988 1991 1940 1999 1995a 1995b 1987 1988a 1988b 1988c 1989a 1989b 1989c 1990
1991a 1991b 1991c 1992a 1992b 1992c 1992d 1993a 1993b 1993c 1993d 1993e 1993f
1993g 1993h 1994a 1994b 1994c 1994d 1996 1998a 1998b 1994 1979 1982 1992 1993
1974 1998a 1998b 1999 1996 1999 1977 1939 1940a 1940b 1961 1990 1991 1994 1990
1991 1996a 1996b 1977 1992 1992 1991 1973 1980 1975 1987 1986 1993a 1993b 1994
1984 1989 1985a 1985b 1987 1987 1986 1992 1994 1986 1987 1991 1990 1995 1997 1995
1993 1990 1985 1990a 1990b 1989 1995 1990 1991 1995 1996 1971 1991 1996 1972 1983
1988a 1988b 1993 1994a 1994b 1994c 1995 1991 1992 1996a 1996b 1993a 1993b 1995a
1995b 1999 1986 1991 1977 1978 1981 1998 1977 1992 1985 1998 1992 1993 1993 1985
1990 1986 1987 1994 1986 1987 1988 1989a 1989b 1991 1992 1993a 1993b 1993c 1994
1996a 1996b 1998 1989 1996 1989 1991 1984 1986 1984 1993 1996 1991 1962 1990
1997 1982 1938 1956 1966 1973 1997 1992 1994 1989 1990 1989 1980 1964 1991 1997
1987 1990 1988 1987 1993 1996 1988 1996 1970 1982 1986 1987a 1987b 1989 1984
1994 1998 1968 1998 1987 1986 1992 1990 1991 1989 1984 1980 1992 1992 1992 1993a
1993b 1995 1996a 1996b 1986 1987 1977 1972 1974 1987 1990 1989 1993 1990 1992
1985 1996 1995 1990 1987 1995 1993 1974 1984a 1984b 1987a 1987b 1983 1990 1992
19 1987 1967 1995 1997 1989 1994 1992 1987a 1987b 1988 1980 1984 1986 1987 1987
1989 1990 1993 1988 1984 1981 1982 1973 1982 1981 1989 1982 1973a 1973b 1974 1970
1987 1987 1997 1996 1995 1977 1993 1980 1987 1953 1985 1991 1992 1998 1997 1996
1981 1993 1987 1994 1987 1998 1987 1989 1995 1996 1970 1971a 1971b 1971c 1971d