

CHAIN-LEVEL MODELS OF EQUIVARIANT TOPOLOGICAL HOCHSCHILD HOMOLOGY OF SEMIPERFECT AND SMOOTH ALGEBRAS OVER FINITE FIELDS

VIKTOR BURGHARDT, PO HU, DANIEL KRIZ, IGOR KRIZ AND PETR
SOMBERG

ABSTRACT. We find an explicit model for equivariant topological Hochschild homology of semiperfect and smooth algebras over a finite field, as modules over the constant Green functor over a primary cyclic group.

1. INTRODUCTION

In [3], Bhatt, Morrow, and Scholze used the fixed point spectrum of equivariant topological Hochschild homology to define the motivic filtration on the De Rham-Witt complex of a smooth \mathbb{F}_p -algebra (first considered by Nygaard [23]). A key new idea in their approach was faithfully flat descent to quasiregular semiperfect \mathbb{F}_p -algebras. On the one hand, this was an application of topological Hochschild homology to obtaining a new technique in p -adic Hodge theory, on the other hand, it gave spectral sequences linking crystalline cohomology to different variants of topological cyclic homology.

An analogous story is also developed in [3] in mixed characteristic, where one can similarly treat quasisyntomic rings using descent to quasiregular semiperfectoid \mathbb{Z}_p -algebras. The role of crystalline cohomology is then replaced by the new concept of prismatic cohomology. In addition to further advancing the methods of p -adic Hodge theory, the spectral sequences based on prismatic cohomology led to substantial

1991 *Mathematics Subject Classification.* 13D03, 55N91, 14F30, 74E15.

Key words and phrases. Topological Hochschild homology, Equivariant homology, Mackey functors, Crystalline cohomology, De Rham-Witt complex, motivic filtration.

The authors acknowledge support by grant GA CR 19-28628X. Po Hu acknowledges the support from NSF grant DMS-2301520. Daniel Kriz was supported by PSR grant 2025 - Linea 8 Sottomisura A. Igor Kriz also acknowledges the support of Simons Foundation award 958219.

progress in the calculation of algebraic K-theory (see Antieau, Krause, and Nikolaus [2], also Mathew [21]).

It is a natural question to ask what role topology intrinsically plays in these new constructions. The purpose of this note is to address this question in the case of pure characteristic p , i.e. the case of smooth \mathbb{F}_p -algebras. In this case, it turns out that topology is not involved in the sense that the genuine \mathbb{Z}/p^r -equivariant spectrum

$$(1) \quad TR^{r+1}(R) = THH_{\mathbb{Z}/p^r}(R)$$

can be explicitly modeled by purely algebraic, chain-level objects, i.e. in terms of homological algebra. The main result of our article is to construct and describe such a model. From one point of view, this can be considered a refinement of the spectral sequence method, which only gives associated graded information.

Before describing this in more detail, let us briefly reflect on the category in which we will be doing homological algebra. This is completely answered by genuine equivariant stable homotopy theory, the concepts of which remain in the background of topological Hochschild homology, even though Nikolaus and Scholze [22] used the ‘‘Tate square’’ to largely eliminate genuine equivariant spectra from the discussion. In equivariant stable homotopy theory, generalized Eilenberg-Mac Lane spectra can be represented by chain complexes of Mackey functors. In fact, the (unbounded) derived categories of generalized equivariant Eilenberg-Mac Lane spectra and Mackey functors are equivalent by the work of Greenlees and Shipley [11].

However, in our case, the ground category can be further refined. The reason we are dealing with equivariant generalized Eilenberg-Mac Lane spectra is the calculation of Hesselholt and Madsen [15] proving that the \mathbb{Z}/p^r -equivariant spectrum $TR(\mathbb{F}_p)$ is the Eilenberg-Mac Lane spectrum $H\mathbb{Z}_p$ of the constant Mackey functor \mathbb{Z}_p . This suggests, the right category to be the derived category of E_∞ - $H\mathbb{Z}_p$ -modules, which is equivalent to the (unbounded) derived category of \mathbb{Z}/p^r -Mackey modules over the constant Green functor \mathbb{Z}_p .

It is this category in which we give our chain-level model. Using faithfully flat descent, it suffices to describe the model for quasiregular semiperfect \mathbb{F}_p -algebras. In this case, the model turns out to be quite simple, (thus conforming with the pattern of [3]). The key ingredient is the ‘‘geometric fixed point functor’’ $\Phi^{\mathbb{Z}/p^i}$ from the derived category of \mathbb{Z}/p^s - \mathbb{Z}_p -modules to the derived category of \mathbb{Z}/p^r - \mathbb{Z}_p -modules for $r < s \leq r + i$. In terms of these building blocks, we can state our main result in broad terms as follows:

1. Theorem. *There is an explicit functorial model of $THH_{\mathbb{Z}/p^r}(R)$ for a quasiregular semiperfect \mathbb{F}_p -algebra R in the derived category of \mathbb{Z}/p^r - $\underline{\mathbb{Z}}_p$ -modules which is a direct sum of even suspensions of chain complexes of the form $\Phi^{\mathbb{Z}/p^i}(\underline{\mathbb{Z}}_p)_{\mathbb{Z}/p^s}$.*

A more precise version of this statement is given in Theorem 6 below.

It should be noted that from the point of view of homotopy theory, the cases of positive and mixed characteristic are quite different. While, as already remarked, for \mathbb{F}_p -algebras R , (1) is an E_∞ -module spectrum over $H\underline{\mathbb{Z}}_p$, which enables a purely chain-level model in the derived category of Mackey modules over $\underline{\mathbb{Z}}_p$, for \mathbb{Z}_p -algebras R , this approach only takes us to E_∞ -modules over the \mathbb{Z}/p^r -equivariant spectrum $TR(\mathbb{Z}_p)$, which is known by Bökstedt and Madsen [5] to be a certain form of connective K-theory. Accordingly, the Bott periodicity element occurs in the calculations (see e.g. [2, 5, 21]). Thus, modules over connective K-theory would have to be used to discuss the mixed characteristic case. There is therefore no hope to model (1) in the mixed characteristic case in the derived category of an abelian category. One could hope for a concrete description in terms of some form of connective equivariant (topological) K-theory. This will be left for future work.

For this reason, we stick to the case of pure characteristic p in the present paper. In this case, remarkably enough, with the exception of quasiregular semiperfect descent, we are able to use, more or less, the classical method of Hesselholt-Madsen [15] who calculated the answer for a perfect commutative \mathbb{F}_p -algebra. As already noted, we observe that in this case, (1) is essentially given by the “geometric fixed points” of the constant $H\underline{\mathbb{Z}}_p$ -module over a primary cyclic group \mathbb{Z}/p^s with respect to the subgroup \mathbb{Z}/p^{s-r} (Theorem 3 below). A key observation is that this does not depend on the choice of $s > r$. Explicit chain-level $\underline{\mathbb{Z}}_p$ -Mackey module models are immediately visible. Additionally, geometric fixed points of (1) are also obtained as geometric fixed points of $H\underline{\mathbb{Z}}_p$ with respect to different subgroups.

The other key point is that this result also extends to the case of (1) for quasiregular semiperfect \mathbb{F}_p -algebras R , i.e. that for such R , (1) is described as a direct sum of the same building blocks (i.e. geometric fixed points of the \mathbb{Z}/p^s -equivariant $H\underline{\mathbb{Z}}_p$ with respect to different subgroups). This can also be proved using the methods of [15], along with the Quillen spectral sequence [24], Theorem 5.1 to start the induction. However, to use the results for the program we set out, it is important

to discuss the precise functoriality for the case of quasiregular semiperfect \mathbb{F}_p -algebras. For this purpose, we use another device, namely the module-valued Witt vectors defined by Dotto, Krause, Nikolaus, and Patchkoria [6, 7]. Applying that construction to powers of the augmentation ideal J of the limit perfection S_R of a quasiregular semiperfect \mathbb{F}_p -algebra R , we get subobjects of $W(S_R)$, which can be used to describe our answer functorially (Theorem 6). Additionally, to show that our functoriality is the right one (thereby describing the correct presheaf on the crystalline site), we use a certain rigidity of the geometric fixed points of $H\mathbb{Z}_p$ (Lemma 4).

It is worth noting that other approaches to the motivic filtration are possible, see e.g. Hahn, Raksit, and Wilson [12]. This is related to Hesselholt's description of topological cyclic homology of an associative \mathbb{F}_p -algebra as a derived functor [13]. The latter result is based on a description of TR for a unital tensor \mathbb{F}_p -algebra as a non-commutative analogue of the De Rham-Witt complex. This is then concentrated in homological degrees 0, 1. However, it is worth noting that (1) for a unital tensor \mathbb{F}_p -algebra (or of a smooth commutative \mathbb{F}_p -algebra) is actually not a direct sum of our building blocks given by geometric fixed points of homology with constant coefficients. Rather, finite $RO(\mathbb{Z}/p^r)$ -graded suspensions are involved, which gives non-trivial extensions of the building blocks. This could make discussion of functoriality using this approach substantially more difficult. Therefore, our discussion could be considered as another application of the Bhatt-Morrow-Scholze approach, beyond the motivic filtration.

The present paper is organized as follows: In Section 2, we discuss some basic notation, and recall the theory of Mackey functors over a primary cyclic group, and Mackey modules over the constant Green functor. In Section 3, we introduce our building blocks, and discuss the chain-level model of (1) for a perfect \mathbb{F}_p -algebra R . In Section 4, we discuss the chain-level model for quasiregular semiperfect \mathbb{F}_p -algebras, and the functoriality.

2. PRELIMINARIES

Equivariant stable homotopy theory for a finite group G is the basic background of our discussion. In particular, we are interested in genuine equivariant spectra, which represent $RO(G)$ -graded equivariant homology and cohomology theories. The basic background reference

for this topic is Lewis, May, Steinberger [19]. A G -equivariant spectrum is often decorated by a subscript $E = E_G$, while E^G denotes the corresponding fixed point spectrum. For a genuine equivariant spectrum E and a fixed $n \in \mathbb{Z}$, the system of groups $(\pi_n(E^H))_H$ over subgroups $H \subseteq G$ forms an additive functor from the stable orbit category to abelian groups. Such a functor is known as a *Mackey functor*, which can be described purely algebraically (see [8]). Additionally, there exist *equivariant Eilenberg-Mac Lane spectra* HM for a given Mackey functor M , where $(\pi_n(HM^H))_H$ is equal to M for $n = 0$, and equal to 0 else ([20]).

There is a natural tensor product \square in the category of Mackey functors (see [8, 18]). One can further consider (commutative) ring objects in the category of Mackey functors, which are called *Green functors*. One defines Mackey modules over a Green functor in the obvious way. For a Green functor A , on the other hand, HA is an E_∞ -ring spectrum, and by the work of Greenlees and Shipley ([11], Section 5), the derived category of E_∞ - HA -modules (see [9] for the relevant background) is equivalent to the unbounded derived category of A -Mackey modules.

Recall that for a finite group G and a normal subgroup H , the *geometric H -fixed points* $\Phi^H E$ of a G -spectrum E is the G/H -equivariant spectrum

$$(E \wedge \widetilde{E\mathcal{F}[H]})^H$$

where $\mathcal{F}[H]$ is the family of subgroups of G not containing H , $E\mathcal{F}$ is the classifying space of a family \mathcal{F} (i.e. a G -CW-complex whose K -fixed points are empty if $K \notin \mathcal{F}$ and contractible if $K \in \mathcal{F}$). The symbol \widetilde{X} denotes the unreduced suspension of a G -space X .

From the point of view of representation theory, for a finite-dimensional real G -representation V , one has

$$S^{\infty V} = \widetilde{E\mathcal{F}[V]}$$

where $\mathcal{F}[V]$ is the family of all subgroups $K \subseteq G$ where

$$V^K \neq 0.$$

The essential point is that in the full generality of an A_∞ ring spectrum R , (1) is a genuine \mathbb{Z}/p^r -equivariant spectrum. This was first proved by Bökstedt, Hsiang, and Madsen [4]. More modern approaches now exist, see for example [1, 17]. Therefore, it is of interest to us to study the above concepts explicitly for the groups $G = \mathbb{Z}/p^r$.

Specifically, let us recall that a \mathbb{Z}/p^r -Mackey functor M can be understood as a sequence of abelian groups

$$M_i, \quad i = 0, \dots, r,$$

where M_i is an $\frac{\mathbb{Z}/p^r}{\mathbb{Z}/p^i}$ -module (the value of the functor M at isotropy \mathbb{Z}/p^i). Further, we are given \mathbb{Z}/p^r -equivariant abelian group homomorphisms

$$r = r_i : M_i \rightarrow M_{i-1}, \quad c = c_i : M_i \rightarrow M_{i+1}$$

(the restriction and corestriction) such that, denoting by γ the generator of the ambient group \mathbb{Z}/p^r , we have

$$(2) \quad r_{i+1} \circ c_i = 1 + \gamma^{p^{r-i-1}} + \dots + \gamma^{p^{r-i-1}(p-1)}.$$

Modules over the \mathbb{Z}/p^r -equivariant Green functor $\underline{\mathbb{Z}}_p$ are sequences of \mathbb{Z}_p -modules M_i , $i = 0, \dots, r$ which satisfy the above axioms and the additional property

$$(3) \quad c_i \circ r_{i+1} = p$$

(see e.g. [18]). Morphisms are tuples of homomorphisms of \mathbb{Z}_p -modules which commute with restrictions and corestrictions.

The principal projectives in the abelian category of \mathbb{Z}/p^r -equivariant $\underline{\mathbb{Z}}_p$ -modules are modules of the form

$$(4) \quad \underline{M}[\mathbb{Z}/p^i]$$

where $0 \leq i \leq r$, M is a projective \mathbb{Z}_p -module and

$$\underline{M}[\mathbb{Z}/p^i]_j = \begin{cases} M[\frac{\mathbb{Z}/p^r}{\mathbb{Z}/p^j}] & \text{for } r - j \leq i \\ M[\frac{\mathbb{Z}/p^r}{\mathbb{Z}/p^{r-i}}] & \text{for } r - j > i. \end{cases}$$

The restrictions and corestrictions are given by

$$r_j(1) = 1 \text{ for } j \leq r - i$$

and

$$c_j(1) = 1 \text{ for } j \geq r - i.$$

(The remaining corestrictions and restrictions are determined.) One can think of (4) as the free \mathbb{Z}/p^r -equivariant $\underline{\mathbb{Z}}_p$ -module freely generated by M in isotropy \mathbb{Z}/p^{r-i} .

3. THE BUILDING BLOCKS AND THE CASE OF PERFECT
 \mathbb{F}_p -ALGEBRAS

Let α_s be the irreducible complex representation of \mathbb{Z}/p^s given by sending the generator γ to $\zeta_{p^s} = e^{2\pi i/p^s}$. Let $\alpha_{s,i} = \alpha_s^{p^i}$ for $i = 0, \dots, s$. Let, in the derived category $D\mathbb{Z}/p^r\text{-Mod}$ of modules over the constant \mathbb{Z}/p^r -Mackey functor $\underline{\mathbb{Z}}_{\mathbb{Z}/p^r}$,

$$(5) \quad \mathcal{W}_{r,j} = (\tilde{C}_*(S^{\infty\alpha_{s,i}}))^{\mathbb{Z}/p^{s-r}}, \quad i \geq 0, \quad i = j - r + s - 1,$$

for $j = 0, \dots, r$. Here \tilde{C}_* denotes the ordinary reduced (say, singular) chain complex, with G -action induced by the action on the space. This always gives a chain complex of $\underline{\mathbb{Z}}_p$ -modules (for a more complete discussion, see [18]).

The definition is ambiguous, but by construction, $\mathcal{W}_{r,j}$ is an E_∞ -algebra over $\underline{\mathbb{Z}}_{\mathbb{Z}/p^r}$, whose homotopy type only depends on r and j .

Let us recall that (5) are chain-level models of constructions familiar in equivariant stable homotopy theory. In our present situation, we have

$$(\alpha_{s,i})^{\mathbb{Z}/p^j} = 0 \text{ if and only if } j > i,$$

so

$$S^{\infty\alpha_{s,i}} = E\mathcal{F}[\widetilde{\mathbb{Z}/p^{i+1}}].$$

Taking fixed points on the level of chain complexes and spectra is equivalent. The reason these complexes show up, and the fact that the definition (5) does not depend on s follows from the cyclotomic property of THH , and from the fact that $TR(\mathbb{F}_p) = H\underline{\mathbb{Z}}_p$.

In more detail, TR is the homotopy inverse limit of $THH^{\mathbb{Z}/p^r}$ with respect to the restriction maps (see [4, 15]). This also automatically becomes a genuine \mathbb{Z}/p^r -equivariant spectrum by the isomorphism

$$S^1/(\mathbb{Z}/p^r) \cong S^1.$$

Further, in this sense,

$$TR(\mathbb{F}_p) = H\underline{\mathbb{Z}}_p$$

(which is proved in [15]). We then obtain a map from the geometric fixed points of $TR_{\mathbb{Z}/p^s}$ to the geometric fixed points of $THH_{\mathbb{Z}/p^s}$, which is an equivalence by an immediate computation. Further, the geometric fixed points of THH is THH again by the cyclotomic property.

Further, by the work of Greenlees and Shipley [11], for a Green functor A , the derived category of HA -modules is equivalent to the unbounded derived functors of A -Mackey modules.

By these considerations, we also have a canonical E_∞ -map

$$\pi : \mathcal{W}_{r,j} \rightarrow \mathcal{W}_{r,j'}, \quad 0 \leq j \leq j' \leq r.$$

Comment: One can show that these algebras cannot in general be modeled by strictly (anti-)commutative DGA's. This is because the canonical morphism to the corresponding Tate complex induces an isomorphism on homology groups in non-negative degrees. On the other hand, the Tate complex inherits Steenrod operations from the Borel cohomology complex, which computes the cohomology of a lens space. Therefore, the Tate complex has non-trivial Steenrod operations which restrict to non-trivial Dyer-Lashof operations on the geometric fixed point complex (5).

If we relax our requirements on the coherence of the multiplicative structure, we have the following explicit chain models, where γ denotes a fixed generator of \mathbb{Z}/p^r :

$$(6) \quad \mathcal{W}_{r,0} : \mathbb{Z} \xleftarrow{p^\epsilon} \mathbb{Z}[\mathbb{Z}/p^r] \xleftarrow{1-\gamma} \mathbb{Z}[\mathbb{Z}/p^r] \xleftarrow{p^N} \mathbb{Z}[\mathbb{Z}/p^r] \dots$$

where ϵ is the augmentation and

$$N = 1 + \gamma + \dots + \gamma^{p^r-1},$$

while

$$(7) \quad \mathcal{W}_{r,j} : \mathbb{Z} \xleftarrow{\epsilon} \mathbb{Z}[\mathbb{Z}/p^{r-j+1}] \xleftarrow{1-\gamma} \mathbb{Z}[\mathbb{Z}/p^{r-j+1}] \xleftarrow{N_{r-j+1}} \mathbb{Z}[\mathbb{Z}/p^{r-j+1}] \dots$$

where

$$N_i = 1 + \gamma + \dots + \gamma^{p^i-1}.$$

For any \mathbb{F}_p -algebra A , we have a $(\mathbb{Z}_p)_{\mathbb{Z}/p^r}$ -Mackey module $\underline{W}_{r+1}(A)$ which is $W_{i+1}(A)$ at isotropy \mathbb{Z}/p^i and restrictions resp. corestrictions are given by the Frobenius resp. Verschiebung.

More generally, we also have, for $0 \leq j \leq r$, a $(\mathbb{Z}_p)_{\mathbb{Z}/p^r}$ -Mackey module $\Phi_j \underline{W}_{r+1-j}(A)$ which is $W_{i+1-j}(A)$ at isotropy \mathbb{Z}/p^i for $i \geq j$ and 0 for lower isotropy, and restrictions resp. corestrictions are given by the Frobenius resp. Verschiebung.

2. Lemma. *The \mathbb{Z}/p^r -Mackey homology of $\mathcal{W}_{r,j}$ is given by*

$$(8) \quad H_s(\mathcal{W}_{r,j}) = \begin{cases} \Phi_j \underline{W}_{r+1-j}(\mathbb{F}_p) & \text{for } s \geq 0 \text{ even} \\ 0 & \text{else.} \end{cases}$$

Proof. Use (6), (7). □

Comment: Despite the fact that the homology computed in Lemma 2 consists of $\underline{\mathbb{Z}/p^{(r-j+1)}}$ -modules, it turns out that $\mathcal{W}_{r,j}$ is not in general equivalent to a chain complex in the category of $\underline{\mathbb{Z}/p^{(r-j+1)}}$ -modules.

For example for $r = j = 1$, we have a $\underline{\mathbb{Z}/p}$ -module

$$(9) \quad \mathcal{W}_{1,1} \otimes_{\mathbb{Z}} \underline{\mathbb{Z}/p}.$$

Attempting to construct $\mathcal{W}_{1,1}$ using Bockstein obstruction theory, we get a non-zero obstruction at the $(p-1)$ 'st step, indicating a non-zero higher Massey power of the generator of an odd-degree cohomology group (one can also see this effect on the Tate complex, where the computation is simpler).

Nevertheless, we have the following:

3. Theorem. *Let R be a perfect \mathbb{F}_p -algebra. Then in the category of Mackey modules $D\underline{\mathbb{Z}/p}$ -Mod, one has*

$$(10) \quad THH_{\underline{\mathbb{Z}/p^r}}(R) = \mathcal{W}_{r,0} \otimes_{\mathbb{Z}} W(R).$$

Proof. The proof is a complete rehash of the case of $R = \mathbb{F}_p$ treated by Hesselholt and Madsen [15]. We use induction on r using the cyclotomic property, and determining the differential in the Borel homology spectral sequence, which follows from the Tate spectral sequence. This also gives the identification of the connecting map and, by induction the $\underline{\mathbb{Z}/p}$ -module model of $THH_{\underline{\mathbb{Z}/p^r}}(R)$. \square

It is also useful to record the morphisms between the objects $\mathcal{W}_{r,j}$ of the (unbounded) derived category of $\mathcal{W}_{r,0}$ -modules, which can be thought of as a partial “rigidity” statement:

4. Lemma. *The graded module of morphisms from $\mathcal{W}_{r,i}$ to $\mathcal{W}_{r,j}$ in $D\mathcal{W}_{r,0}$ is isomorphic to the \mathbb{Z}/p^r isotropy part of $H_*\mathcal{W}_{r,j}$ when $i \leq j$ and is 0 else.*

Proof. Recall the \mathbb{Z}/p^s -equivariant based CW-complexes $\Phi_{s,i}$ $0 < i \leq s$ which are equivalent to S^0 in isotropies \mathbb{Z}/p^j for $j \geq i$ and contractible in smaller isotropies. Then the suspension spectra $\Sigma^\infty \Phi_{s,j}$ are E_∞ -ring spectra, where

$$(11) \quad \Phi_{s,i} \wedge \Phi_{s,j} \sim \Phi_{s,\max(i,j)}.$$

The groups in question can be expressed as derived \mathbb{Z}/p^r -equivariant maps between $\mathcal{W}_{r,0}$ -module of the form $\mathcal{W}_{r,0} \wedge \Phi_{s,i}$ for different i . Thus, the positive part of our statement (the case of $i \leq j$) follows from (11).

For the case $i > j$, consider a map of $\mathcal{W}_{r,0}$ -modules

$$(12) \quad \Sigma^n \mathcal{W}_{r,i} \rightarrow \mathcal{W}_{r,j}.$$

Composing with the n -suspension of the projection

$$(13) \quad \mathcal{W}_{r,j} \rightarrow \mathcal{W}_{r,i},$$

we obtain a map of $\mathcal{W}_{r,0}$ -modules

$$(14) \quad \Sigma^n \mathcal{W}_{r,j} \rightarrow \mathcal{W}_{r,j},$$

which we already classified. In particular, we claim that the composition (14) is 0. Otherwise, it would be injective on homotopy groups. However, we also know that the projection (13) is 0 on homotopy groups in degrees $n \gg 0$.

Thus (14) is 0. Thus, the map (12) factors through the cofiber C of the projection (13). However, there are no non-zero morphisms of $H\mathbb{Z}$ -modules

$$(15) \quad \Sigma^n C \rightarrow \mathcal{W}_{r,j},$$

since C has a $H\mathbb{Z}$ -module cellular structure where cells are principal projectives with generators in isotropies in which $\mathcal{W}_{r,j}$ has 0 homotopy (see (7)).

However, then there are also no non-zero morphisms (15) of $\mathcal{W}_{r,0}$ -modules, since

$$C \wedge_{H\mathbb{Z}} \mathcal{W}_{r,0} \sim C.$$

□

4. THE CASE OF A SEMIPERFECT \mathbb{F}_p -ALGEBRA

We now treat the case of a quasiregular semiperfect \mathbb{F}_p -algebra R . We will give the definition of this concept, which is somewhat technical, following [3], but for our purposes, we can focus on the case

$$(16) \quad R = C \otimes_A C \otimes_A \cdots \otimes_A C$$

where A is a smooth \mathbb{F}_p -algebra and C is its colimit perfection.

The reason for this is that if A is a smooth \mathbb{F}_p -algebra, we can consider cosimplicial descent from A to (16). A crucial fact used in [3] is that this preserves (1). Therefore, to give a \mathbb{Z}_p -module chain model of $THH(A)$ for a smooth \mathbb{F}_p -algebra A , it suffices to give it for quasiregular semiperfect algebras, specifically those of the form (16).

Denote by

$$S_R = \lim(\dots \xrightarrow{\phi} R \xrightarrow{\phi} R)$$

the limit perfection of R , where $\phi : R \rightarrow R$ is the Frobenius.

Denote by J the kernel of the projection $S_R \rightarrow R$ and denote also

$$I = \text{Ker}(\phi : R \rightarrow R).$$

One has

$$J/J^2 \cong I/I^2,$$

since $\phi(I) \subseteq I^2$. Using the notation $L_{B/A}$ of Quillen [24] for the derived cotangent complex for a (non-derived) commutative A -algebra B , we have

$$L_{S_R/\mathbb{F}_p} = 0,$$

which, by Theorem 5.1 of [24], gives

$$L_{R/\mathbb{F}_p} = L_{R/S_R}.$$

Then Theorem 6.3 of [24] gives

$$H_1 L_{R/S_R} = \text{Tor}_{S_R}^1(R, R) = J/J^2,$$

which gives

$$(17) \quad H_1 L_{R/\mathbb{F}_p} = I/I^2.$$

By definition ([3], Definition 8.8), a semiperfect \mathbb{F}_p -algebra R is called *quasiregular* when I/I^2 is a flat R -module and

$$H_n L_{R/\mathbb{F}_p} = 0 \text{ for } n \neq 1.$$

In our situation of interest, we have a somewhat stronger property. Call a quasiregular semiperfect \mathbb{F}_p -algebra R *quasismooth* when J/J^2 is a free R -module of finite rank and

$$(18) \quad \text{Sym}_R^\ell(J/J^2) = J^\ell/J^{\ell+1}.$$

5. Lemma. *Let A be a smooth \mathbb{F}_p -algebra and let P be its colimit perfection. Then*

$$R = P \otimes_A \cdots \otimes_A P$$

is a quasismooth semiperfect \mathbb{F}_p -algebra.

Proof. A tensor product over \mathbb{F}_p of two perfect \mathbb{F}_p -algebras is perfect (since $\phi \otimes \phi$ is a composition of the injective maps $\phi \otimes 1$ and $1 \otimes \phi$). Then one can prove the analogues of the statements for the ideal J for the kernel of the projection $P \rightarrow R$ (since this situation is a colimit of the smooth case, where the statement follows from the theory of parameters). Now while P is different from the limit perfection S_R , one can also prove that the statement about the ideal J is equivalent

for the kernel ideal of any surjective homomorphism from a perfect ring. \square

Now let R be a quasismooth \mathbb{F}_p -algebra, and let J/J^2 be a free R -module on a finite basis B

$$(19) \quad J/J^2 = \bigoplus_B R.$$

We shall write

$$(20) \quad W_r^{B_\ell}(R) = \bigoplus_{B_\ell} W_r(R)$$

where B_ℓ is the set of unordered ℓ -tuples (with possible repeats) of elements of B .

We can write this construction more functorially and more generally as follows. Recalling the notion of Witt vectors with coefficients in a module [6, 7, 25], we have

$$(21) \quad W_r^{B_\ell}(R) \cong W_r(S_R, J^\ell)/W_r(S_R, J^{\ell+1}).$$

We see that the right-hand side does not depend on the choice of a basis, and, in fact, makes sense for every quasiregular semiperfect \mathbb{F}_p -algebra R . In that generality, this motivates in fact writing

$$\mathcal{W}_{r,j}(R, \ell) := W(S_R, J^\ell)/W(S_R, J^{\ell+1}) \otimes \mathcal{W}_{r,j}.$$

Thus, we may also write

$$\mathcal{W}_r(R, \ell) := \text{holim } \mathcal{D}_r(R, \ell),$$

where $\mathcal{D}_r(R, \ell)$ is the diagram

$$\begin{array}{ccc} & & \mathcal{W}_{r,r}(R, \ell) \\ & & \downarrow \phi \\ & \mathcal{W}_{r,r-1}(R, \ell) \xrightarrow{\pi} & \mathcal{W}_{r,r}(R, \ell) \\ & \downarrow \phi & \\ \dots & \xrightarrow{\pi} & \mathcal{W}_{r,r-1}(R, \ell) \\ & \downarrow \phi & \\ \mathcal{W}_{r,0}(R, \ell) \xrightarrow{\pi} & & \dots \end{array}$$

where ϕ is the map induced by the p -th power (i.e. Frobenius) on R .

6. Theorem. *For a quasiregular semiperfect \mathbb{F}_p -algebra R , one has*

$$(22) \quad THH^{\mathbb{Z}/p^r}(R)_{2\ell} = \bigoplus_{i=0}^{\ell} W_{r+1}(S_R, J^i)/W_{r+1}(S_R, J^{i+1}) \otimes_{W_{r+1}(S_R)} W_{r+1}(R),$$

$$THH^{\mathbb{Z}/p^r}(R)_{2\ell+1} = 0.$$

Further, in the category $D\mathbb{Z}_p\text{-Mod}$, one has

$$(23) \quad THH_{\mathbb{Z}/p^r}(R) = \bigoplus_{\ell \geq 0} \mathcal{W}_r(R, \ell)[2\ell].$$

7. Corollary. *If R is a quasiregular semiperfect \mathbb{F}_p -algebra, then $THH_{\mathbb{Z}/p^r}(R)$ is equivalent to a direct sum of even suspensions of the \mathbb{Z}_p -module complexes $\mathcal{W}_{r,i}$.*

Proof of Theorem 6. We first consider the case $r = 0$. Then by [24], Theorem 5.1, applied to the sequence

$$\mathbb{F}_p \rightarrow R \otimes R \rightarrow R,$$

we obtain

$$L_{R/R \otimes R} = J/J^2[2].$$

By [24], Theorem 6.3, we then have

$$\begin{aligned} Tor_{2\ell}^{R \otimes R}(R, R) &= Sym_R^\ell J/J^2 = J^\ell/J^{\ell+1}, \\ Tor_{2\ell+1}^{R \otimes R}(R, R) &= 0. \end{aligned}$$

Now following [15], we have a spectral sequence

$$Tor_*^{A_* \otimes R \otimes R}(R, R) \Rightarrow THH_*(R)$$

where A_* is the dual Steenrod algebra which, using the standard differentials of [15] for $p > 2$, implies (22) (and hence (23)) for $r = 0$.

For $r \geq 1$, one then also repeats the method of [15], considering the Borel homology spectral sequence, which has the differential induced from the case $R = \mathbb{F}_p$ and consequently collapses to even degrees. Extensions are given by multiplication by p composed with the map induced by the Frobenius on R .

One then again gets the total fixed points by using the fundamental cofibration

$$E\mathbb{Z}/p_+^r \wedge E \rightarrow E \rightarrow \widetilde{E\mathbb{Z}/p} \wedge E$$

for a \mathbb{Z}/p^r -equivariant spectrum E , and identifying the last term by induction using the cyclotomic property. The identification of the connecting map also mimics the perfect case, thus giving the result as well as the decomposition (23).

The induction described is sufficient to prove the statement of Corollary 7, and to count the number of copies of the complexes $\mathcal{W}_{r,i}$. However, we stated our answer functorially in R , and the functoriality must be right in the A_∞ -sense to be used for descent. For this purpose, we invoke Lemma 4. We first observe that in the limit, we have

$$(24) \quad TF_{\mathbb{Z}/p^r}(R) \sim \bigoplus_{\ell \geq 0} \underline{W(S_R, J^\ell/J^{\ell+1})}[2\ell].$$

Since $W(S_R, J^\ell/J^{\ell+1})$ are free \mathbb{Z}_p -modules, there are no higher derived maps between the \mathbb{Z}_p -modules (24). Now let $R \rightarrow R'$ be a homomorphism of quasiregular semiperfect \mathbb{F}_p -algebras. Then the functoriality we describe fits into the diagram of $H\mathbb{Z}_p$ -modules

$$(25) \quad \begin{array}{ccc} TF_{\mathbb{Z}/p^r}(R) & \longrightarrow & TF_{\mathbb{Z}/p^r}(R') \\ \downarrow & & \downarrow \\ THH_{\mathbb{Z}/p^r}(R) & \longrightarrow & THH_{\mathbb{Z}/p^r}(R') \end{array}$$

where the horizontal arrows are given by our functoriality, while the vertical arrows are the natural projections. We may extend scalars over $H\mathbb{Z}_p = TF_{\mathbb{Z}/p^r}(\mathbb{F}_p)$ to $THH_{\mathbb{Z}/p^r}(\mathbb{F}_p)$ to obtain a diagram of $\mathcal{W}_{r,0}$ -modules

$$(26) \quad \begin{array}{ccc} TF_{\mathbb{Z}/p^r}(R) \otimes_{\mathbb{Z}_p} \mathcal{W}_{r,0} & \longrightarrow & TF_{\mathbb{Z}/p^r}(R') \otimes_{\mathbb{Z}_p} \mathcal{W}_{r,0} \\ \downarrow & & \downarrow \\ THH_{\mathbb{Z}/p^r}(R) & \longrightarrow & THH_{\mathbb{Z}/p^r}(R'). \end{array}$$

By Lemma 4, however, the bottom horizontal arrow (25) is uniquely determined (in the A_∞ -sense) by the remaining arrows. This completes the proof of our functoriality statement. \square

Comment: Theorem 6 can be thought of as specifying an A_∞ -sheaf on the crystalline site of \mathbb{F}_p -algebras considered in [3]. Therefore, we can use faithfully flat quasiregular semiperfect descent to obtain Mackey chain models for $THH_{\mathbb{Z}/p^r}(R)$ for a smooth \mathbb{F}_p -algebra R .

The ‘‘rigidity’’ of Lemma 4, however, does not imply that in the case of smooth \mathbb{F}_p -algebras, we would get again sums of copies of the complexes $\mathcal{W}_{r,i}$. In fact, in the case of the polynomial algebra $\mathbb{F}_p[x]$, it is known ([10, 13, 14, 16]) that one obtains summands of finite $RO(\mathbb{Z}/p^r)$ -graded suspensions of the complexes $\mathcal{W}_{r,i}$, which are non-trivial extensions of the complexes $\mathcal{W}_{r,i}$. In the case of quasiregular semiperfect

algebras, this behavior is ruled out in part by the fact that the chain homology is contained in even degrees.

However, it is worth noting that by generalizing the Illusie complex to tensor algebras, Hesselholt [13] was able to describe TC of an \mathbb{F}_p -algebra as a derived functor. Hahn, Raksit, and Wilson [12] recently used a similar alternative to quasiregular semiperfect descent to define a version of the motivic filtration on commutative ring spectra.

REFERENCES

- [1] V. Angeltveit, A. J. Blumberg, T. Gerhardt, M. A. Hill, T. Lawson, M. Mandell: Topological cyclic homology via the norm, *Doc. Math.* 23 (2018), 2101-2163
- [2] B. Antieau, A. Krause, T. Nikolaus: On the K-theory of \mathbb{Z}/p^n , arXiv: 2405.04329
- [3] B. Bhatt, M. Morrow, P. Scholze: Topological Hochschild homology and integral p -adic Hodge theory, *Publ. Math. Inst. Hautes Études Sci.* 129 (2019), 199-301
- [4] M. Bökstedt, W. C. Hsiang, I. Madsen: The cyclotomic trace and algebraic K-theory of spaces, *Invent. Math.* 111 (1993), no. 3, 465-539
- [5] M. Bökstedt, I. Madsen: Topological cyclic homology of the integers, *Astérisque*, v. 226 (1994), p. 57-143
- [6] E. Dotto, A. Krause, T. Nikolaus, I. Patchkoria: Witt vectors with coefficients and characteristic polynomials over non-commutative rings, arXiv: 2002.01538
- [7] E. Dotto, A. Krause, T. Nikolaus, I. Patchkoria: Witt vectors with coefficients and TR, arXiv: 2312.12971
- [8] A. W. M. Dress: Contributions to the theory of induced representations *Lecture Notes in Mathematics* 342, (1973), 183-240
- [9] A. Elmendorf, I. Kriz, M. Mandell, P. May: *Rings, modules and algebras in stable homotopy theory*, AMS Mathematical Surveys and Monographs, Volume 47, 1997
- [10] T. Gerhardt: The $RO(S^1)$ -graded equivariant homotopy of $THH(\mathbb{F}_p)$, PhD thesis, MIT, 2007
- [11] J. P. C. Greenlees and Brooke Shipley: Fixed point adjunctions for equivariant module spectra, *Algebr. Geom. Topol.* 14 (2014) 1779-1799
- [12] J. Hahn, A. Raksit, D. Wilson: Motivic filtration on the topological cyclic homology of commutative ring spectra, to appear in *Annals of Mathematics*, arXiv:2206.11208v2
- [13] L. Hesselholt: Witt vectors of non-commutative rings and topological cyclic homology, *Acta Math.* 178 (1997), no. 1, 109-141, Correction: *Acta Math.* 195 (2005), 55-60
- [14] L. Hesselholt: K-theory of truncated polynomial algebras, *Handbook of K-theory*. Vol. 1, 2, 71-110. Springer-Verlag, Berlin, 2005
- [15] L. Hesselholt, I. Madsen: On the K-theory of finite algebras over Witt vectors of perfect fields, *Topology* 36 (1997), no. 1, 29-101

- [16] P. Hu, I. Kriz, P. Somberg: On the equivariant motivic filtration of the topological Hochschild homology of polynomial algebras, *Adv. Math.* 412 (2023), Paper No. 108803, 12 pp.
- [17] P. Hu, I. Kriz, P. Somberg, F. Zou: Equivariant operations in topological Hochschild homology, *Math. Z.* 310 (2025), no. 4, Paper No. 67, 21 pp.
- [18] S. Kriz: Some Remarks on Mackey Functors, arXiv:2205.12192
- [19] L. G. Lewis, Jr., J. P. May, M. Steinberger, and J. E. McClure: *Equivariant stable homotopy theory*, Vol 1213 of Lecture Notes in Mathematics, Springer-Verlag, Berlin, 1986.
- [20] L. G. Lewis, Jr., J. P. May, M. Steinberger, and J. E. McClure: Ordinary $RO(G)$ -graded cohomology, *Bull. Amer. Math. Soc. (N.S.)* 4 (1981), no. 2, 208-212
- [21] A. Mathew: Some recent advances in topological Hochschild homology, *Bull. Lond. Math. Soc.* 54 (2022), no. 1, 1–44
- [22] T. Nikolaus, P. Scholze: On topological cyclic homology, *Acta Math.* 221 (2018), no. 2, 203-409, correction *Acta Math.* 222 (2019), no. 1, 215-218
- [23] N. O. Nygaard: Slopes of powers of Frobenius on crystalline cohomology, *Ann. Sci. ENS* 14 (1981), 369-401
- [24] D. Quillen: On the (co-) homology of commutative rings, *Applications of Categorical Algebra* (Proc. Sympos. Pure Math., Vol. XVII, New York, 1968), pp. 65-87 Proc. Sympos. Pure Math., XVII American Mathematical Society, Providence, RI, 1970
- [25] T. Read: p -typical Witt vectors with coefficients and the norm, arXiv: 2305.20064